



Physical and Biochemical Evaluation of Blueberries during Long-term Frozen Storage at -20 °C: Implications for Shelf-life and Quality Preservation

Sahar Dahbi^{1*}, Hafida Hanine², Ossama Kodad³, Mohammed Raouane¹, Souad Amghar¹

1 Reseach Team "Lombricidae, Improving Soil Productivity and Environment" (LAPSE), Centre "Eau, Ressources Naturelles, Environnement et Développement Durable" (CERNE2D), Ecole Normale Supérieure (ENS), Mohammed V University in Rabat, Rabat, Morocco"

2 Laboratory of Industrial Engineering and Surface Engineering, Faculty of Science and Techniques, Sultan Moulay Slimane University, Beni Mellal, Morocco

3 Department of Arboriculture-Viticulture, Ecole Nationale d'Agriculture de Meknès, Meknès, Morocco

ARTICLE INFO

*Corresponding author's email: saharenam69@gmail.com

Article history:

Received: 6 December 2023,

Received in revised form: 30 March 2024,

Accepted: 2 April 2024

Article type:

Research paper

Keywords:

Antioxidant activity,
Bioactive compounds,
Blueberry,
Quality preservation,
Shelf-life extension,
Subzero storage

ABSTRACT

Blueberry (*Vaccinium spp.*) is well-known for its high antioxidant capacity and exceptional richness of vitamins, fiber, and polyphenols. Due to its bioactive compounds, flavonoids, tannins, and phenolic acids, blueberry possesses various health benefits, such as anti-diabetic, anti-cardiovascular, and anti-carcinogenic properties. However, its limited production period and short shelf-life generate tremendous post-harvest loss. In this context, post-harvest treatments are necessary to extend the shelf life of blueberries and promote their consumption throughout the year. Thus, this work aimed to evaluate changes in the physical and biochemical quality of blueberries stored at -20 °C for three months. The results showed that one month of frozen storage increased the anthocyanins content in blueberries and preserved all other characteristics. The biochemical fruit quality improved in the second month, while the titratable acidity, pH, total soluble solids, and weight remained preserved. However, at the end of the storage period, there was a continual rise in anthocyanins and flavonoids, whereas polyphenols and antioxidant activity remained stable, surpassing the levels found in fresh blueberries. A notable decline in physical characteristics was observed. In this context, it is recommended to store blueberries at -20 °C for one month at most to maintain fresh consumption quality. Blueberries stored for over two months must be intended for processed blueberry-based products to fully utilize their richness in phenolic compounds and avoid issues related to their sour taste and reduced weight.

Abbreviations: Antioxidant activity (AA), Phenylalanine ammonia-lyase (PAL), Titratable Acidity (TA), Total Anthocyanins Content (TAC), Total Flavonoid content (TFC), Total Polyphenol content (TPC), Total Soluble Solids (TSS)

Introduction

Blueberry (*Vaccinium spp.*) belongs to the

Ericaceae family, specifically to the genus *Vaccinium*, which contains approximately 450

COPYRIGHT

© 2025 The author(s). This is an openaccess article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other medium is permitted, provided the original author(s) and source are cited, in accordance with accepted academic practice. No permission is required from the authors or the publishers.

species (Edger et al., 2022). Since prehistoric times, the blueberry originated from a spontaneous shrub native to the northern regions of Europe and North America. Its distribution expanded in cultivation, regions with mild winters, such as Mediterranean countries (Edger et al., 2022; Pistón et al., 2017). In the last decade, the importance of blueberry species has grown due to their recognized health benefits, including antidiabetic, anti-cardiovascular, and anticancer potential (Silva et al., 2020). These properties are attributable to various bioactive compounds such as flavonoids, tannins, and phenolic acids (Seeram, 2008; Smith et al., 2000). Also, blueberries have a high antioxidant capacity against free radicals and reactive species (Rodríguez-Mateos et al., 2012; Silva et al., 2020; Veberic et al., 2015; Vrhovsek et al., 2012).

However, blueberries are renowned as perishable fruits due to their short shelf life and limited production period, resulting in notable postharvest loss (Duan et al., 2022). Thus, postharvest treatments are necessary to overcome these wastes, including cold storage, heat treatments, irradiation, etc. (Mahajan et al., 2014).

The present work aimed to study the impact of a specific postharvest treatment, specifically the storage of blueberries at -20 °C for three months. The objective was to consider their physical and biochemical characteristics as they changed in storage at the subzero temperature.

Materials and Methods

Plant materials

The plant materials in this study originated from an estate in the Kenitra region in Morocco. The fruits of the “Southern Highbush” cultivar were hand-harvested in March 2017 based on their external appearance: fully colored berries (90%-100%) and absence of visible lesions and rot. Subsequently, the samples were partitioned into three freezer bags, with each containing 42 g of the sample. We stored the samples at -20 °C for three months and took a sample for physicochemical and biochemical analysis at the end of every month.

Phytochemical analysis

Total soluble solids (TSS)

The TSS was measured using a refractometer (ATAGO C.O Ltd; Model PR-1). The results were expressed as °Brix.

pH

Freshly extracted blueberry juice was measured for pH using an electronic pH meter.

Weight

Total weight was counted using a precision electronic scale. The results were expressed in grams.

Titrateable acidity (TA)

The titrateable acidity was carried out based on a protocol described by Nielsen et al. (2017). Five g of freshly extracted juice from each sample was added to 25 mL of distilled water. The process involved 0.1 N (NaOH) to neutralize the acids present in blueberry juice until the pH reached 8.1.

The results were expressed as citric acid (%) (Nielsen, 2017).

$$\% \text{ TA} = \frac{V(\text{NaOH}) \times \text{Conc}(\text{NaOH}) \times K \times 100}{g \text{ of the sample}}$$

Where V (NaOH) is the required volume of NaOH in mL to reach pH 8.1.

Conc (NaOH) is the NaOH concentration.

K is the specific coefficient of the dominant acid in blueberry juice (citric acid) equal to 0,064 (Adi et al., 2019).

Biochemical analysis

Extract

The extraction involved adding five g of blueberries to 25 mL of methanol and a drop of HCl (0.1% hydrochloric acid). The mixture was stirred using an Ultra Turrax for 15 min and then centrifuged for 15 min at 6000 rpm. Subsequently, methanol was evaporated using a rotary evaporator.

Determination of total polyphenol content (TPC)

Total polyphenols were determined using the Folin-Ciocalteu method (Sánchez-Rangel et al., 2013). The extract was diluted 100 times (D100: 9.9 mL of methanol mixed with 0.1 mL of the sample). Then, 0.25 mL of the sample was added to 0.25 mL of Folin-Ciocalteu reagent, 2 mL of distilled water, and 0.25 mL of sodium carbonate (20%). The mixture was vortexed, and after 30 min of incubation in darkness, the absorbance of the samples was measured using a spectrophotometer at 750 nm. A calibration curve with different concentrations of gallic acid was prepared. The total phenolic content was expressed as milligrams of gallic acid equivalent per gram of fresh sample weight (mg GAE g⁻¹ FW).

Determination of antioxidant activity (AA)

The antioxidant activity was evaluated using the DPPH method, based on a procedure described in

previous studies (Brand-Williams et al. 1995; Mensor et al., 2001). A volume of 0.1 mL of the extract was added to 3.9 mL of DPPH solution (6 mg 100 mL⁻¹ in 80% methanol) with a dilution factor of 100 (D100). After incubation in darkness for 30 min at room temperature, absorbance readings were taken at 515 nm using a spectrophotometer.

The inhibition percentage was calculated using the following formula:

$$I\% = \frac{(Ac - As) \times 100}{Ac}$$

Where Ac is the absorbance of the control (DPPH).

As is the absorbance of the sample (DPPH with the added test sample).

A calibration curve of Trolox solutions was used and the results were expressed as mg of Trolox equivalent per gram of each sample (mg TE g⁻¹).

Determination of total flavonoid content (TFC)

The total flavonoid content (TFC) was assessed according to a method defined by Lamaison J. and Carnat (1990). A volume of 1 mL of each sample was added to 1 mL of AlCl₃ (MeOH). The absorbance was read at 510 nm. Rutin generated a calibration curve, and then the results were expressed as milligrams of rutin equivalent per gram of each sample (mg RE g⁻¹) (Lamaison J. and Carnat, 1990).

Determination of total anthocyanins content (TAC)

A procedure for determining anthocyanin content was based on a protocol described by Giusti and Wrolstad (2001). Two buffers were employed, i.e., a potassium chloride buffer at pH 1.0 and a sodium acetate buffer at pH 4.5. Accordingly, 0.4 mL of the diluted extract was separately mixed with 3.6 mL of each of the two buffers, and then the absorbance was measured by a spectrophotometer at 510 nm and 700 nm. (Giusti and Wrolstad, 2003)

The final absorbance value was obtained by subtracting the measured absorbance at different wavelengths.

$$Abs = (Abs_{510} - Abs_{700})_{pH1} - (Abs_{510} - Abs_{700})_{pH4.5}$$

The results were expressed as micrograms of cyanidin-3-glucoside equivalent per gram of each sample (mg CGE g⁻¹) and were calculated using the following equation:

$$TAC \text{ (mg CGE g}^{-1}\text{)} = \frac{Abs \times DF \times MW \times 100}{MAC}$$

Where Abs is absorbance.

DF is dilution factor (100).

MW is the molecular weight of cyanidin-3-glucoside (449,2) (Olivas-Aguirre et al., 2016).

MAC is the molar absorptivity of cyanidin-3-glucoside (26.900) (Young and Abdel-Aal, 2009).

Statistical analysis

All results were statistically conducted using IBM SPSS Statistics 20 software. Both one-way and multi-way analysis of variance (ANOVA) and the Student-Newman-Keuls (SNK) tests were utilized for specific mean comparisons. Statistical significance was considered per treatment group (p<0.05). Three replicates were considered for each analysis, and the results were expressed as mean values ± standard deviation (SD).

Results

Biochemical characteristics

Total polyphenol content (TPC)

Average polyphenol content in the fresh blueberries was 2.631 ± 0.168 mg GAE g⁻¹. After one month of freezing storage at -20 °C, TPC remained statistically stable (p = 0.906), indicating that no polyphenol loss occurred in blueberries stored at -20 °C for 30 days. After two months of cold storage, we noticed a significant increase compared to fresh fruits. After three months of frozen storage, the polyphenol content continued to rise in a non-significant manner (p=0.147), reaching its highest value at 11.366 ± 1.1621 mg GAE g⁻¹ (Fig. 1).

Antioxidant activity

The antioxidant activity in blueberries exhibited a similar trend of changes as observed in polyphenol content. The results showed that during the first month of the storage period, the antioxidant activity remained at the same level (p=0.228). In the second month, the antioxidant activity increased by 54.57% compared to the berries in their initial state. At the end of freezing storage, a highly significant increase (53.99%) was recorded, compared with the initial fresh berries. The value slightly increased in a non-significant manner compared to the previous month (p = 0.468) (Fig. 2).




Fig. 1. Comparative analysis of polyphenol levels in three months of frozen storage. a and b are different letters that indicate significant differences ($p < 0.05$).




Fig. 2. Changes in antioxidant activity during the three months of frozen storage. a, b: different letters indicate significant differences ($p < 0.05$).

Total flavonoid content (TFC)

Changes in total flavonoid content in blueberries stored at $-20\text{ }^{\circ}\text{C}$ showed an interesting trend over time. After one month, blueberries showed a non-significant rise in flavonoids ($p = 0.078$) compared to the fresh samples. However, remarkable changes were detected throughout the latter part of the freezing storage period. A

significant increase occurred at the end of the second month. The flavonoid content increased from $2.46 \pm 0.221\text{ mg RE g}^{-1}$ to $4.47 \pm 0.177\text{ mg RE g}^{-1}$ compared to the previous month. Meanwhile, the flavonoid content of blueberries stored for three months reached a value of $5.172 \pm 0.097\text{ mg RE g}^{-1}$, which denotes a rise of 146% compared to fresh fruits (Fig. 3).




Fig. 3. Variations in total flavonoid content across three months of frozen storage. a, b: different letters indicate significant differences ($p < 0.05$).

Total anthocyanin content (TAC)

The TAC improved significantly by 5% during the first month of frozen storage. However, no significant changes were observed during the

second month. By the end of the frozen storage, a significant increase (7%) occurred compared to the fresh berries, reaching a value of $169.744 \pm 3.200\text{ mg CGE g}^{-1}$ (Fig. 4).




Fig. 4. Changes in anthocyanin levels over three months of frozen storage. a, b: different letters indicate significant differences ($p < 0.05$).

Titrateable acidity (TA) and pH level

As illustrated in Figure 5, the acidity of frozen blueberries increased through the storage period. However, these changes were not significantly

different over the first two months ($p = 0.164$). Conversely, a statistically significant rise was observed at the end of the third month of frozen storage.




Fig. 5. Changes in titrateable acidity over three months of storage at $-20\text{ }^{\circ}\text{C}$. a, b: different letters indicate significant differences ($p < 0.05$).

This increase in sour taste was supported by a decrease in pH value. After one month of frozen storage, the pH of blueberries decreased insignificantly from 3.28 ± 0.014 to 2.81 ± 0.035

(Fig. 6). After three months, a significant decrease of 38% was observed compared to the fresh fruits, reaching a minimum value (2.6 ± 0.060).




Fig. 6. Changes in pH throughout the three months of frozen storage. a, b: different letters indicate significant differences ($p < 0.05$).

Total soluble solids (TSS)

The total soluble solids content in fresh blueberries was 14.5 ± 0.707 °Brix. However, this value decreased by 12% after 30 days of frozen storage ($-20\text{ }^{\circ}\text{C}$) and remained significantly stable throughout the second month. Then, a notable reduction (33%) in soluble solids content was

observed, reaching a minimum of 9.75 ± 0.071 °Brix (Fig. 7).

Weight

According to Figure 8, storing blueberries at $-20\text{ }^{\circ}\text{C}$ significantly preserved their weight until the end of the second month. At the end of the

third month, a significant decrease (13.26%) was observed compared to the fresh fruits.




Fig. 7. Changes in total soluble solids (TSS) during the three months of storage (-20 °C). a, b: different letters indicate significant differences ($p < 0.05$).




Fig. 8. Changes in weight during the frozen storage of three months. a, b: different letters indicate significant differences ($p < 0.05$).

Discussion

Blueberry is a perishable fruit with good health benefits and high economic value. Its beneficial effect emanates from its richness in polyphenolic compounds and antioxidant potential. The total polyphenol content in this research ranged from 2.631 ± 0.168 to 11.366 ± 1.162 mg GAE g^{-1} , which are higher than the values found by Al Hasani et al. (2023). The latter reported an average of 2.09 mg GAE g^{-1} of total polyphenol content (Al Hasani et al., 2023). Indeed, based on the criteria established by Vasco et al. (2008), the researchers classified fruits into three categories having a low (< 1 mg GAE g^{-1}), medium (1–5 mg GAE g^{-1}), or high polyphenol content (> 5 mg GAE g^{-1}) (Vasco et al., 2008). Accordingly, the berries herein are considered a promising source of polyphenolic compounds. Moreover, the fresh blueberries in this research showed a degree Brix of 14.5 ± 0.707 , exceeding the range reported by Matiacevich et al. (2013), which varied between 10.33 and 13.39 depending on the blueberry cultivar (Matiacevich et al., 2013). Nevertheless, it is essential to note that at postharvest, the fruit undergoes a process of senescence and quality deterioration, leading to a rapid loss of its biochemical properties, lower nutrient values, and a shortened shelf life (Tietel et al., 2012). Thus, implementing effective postharvest

treatments such as storage conditions could significantly preserve the biochemical properties of blueberries and, consequently, maintain their health benefits (Tietel et al., 2012; Skrovankova et al., 2015).

In the present research, the study of frozen storage on the quality of blueberries illustrated an increase in biochemical properties, including TPC, AA, TAC, and TFC, by the end of the third month of the storage period. As for the physicochemical characteristics, notable changes occurred after the storage duration, with a rise in titratable acidity. This increase in sour taste was simultaneous with a decrease in pH, TSS, and weight.

Our findings align with some previous studies and reports. For instance, Reque et al. (2014) found that the antioxidant activity in blueberries increased significantly during three months of frozen storage at -18 °C, followed by a decline until the end of six months of storage, while maintaining a value greater than that of the initial (Reque et al., 2014). A parallel pattern occurred in research by Bakowska-Barczak and Kolodziejczyk (2008). The authors noticed an increase in the total anthocyanins in three different Saskatoon berry cultivars stored at -20 °C for nine months. The researchers associated this rise with changes in cyanidin-3-arabinoside in the fruits (Bakowska-Barczak and

Kolodziejczyk, 2008).

Indeed, the increase in bioactive compounds (TPC, AA, TFC, TAC) can stem from reactions that continue to occur in the postharvest stage (Piljac-Žegarac and Šamec, 2011). For example, the increase in phenylalanine ammonia-lyase (PAL) was reportedly due to low-temperature stress (King et al., 2022). Based on reports by other authors, this enzyme (PAL) plays a crucial role in the phenylpropanoid pathway. It activates the production of various beneficial compounds, such as anthocyanins and flavonoids in fruits. These bioactive compounds possess antioxidant properties that assist in preserving nutritional quality. Therefore, PAL potentially enhances berry flavor and health benefits (King et al., 2022; Chen et al., 2006). Another possible explanation could be the degradation of cell structures due to thawing the fruit before analysis, resulting in an easy and abundant extraction of these compounds (De Ancos et al., 2000). Some studies suggest that the increase in anthocyanins might result from the co-pigmentation mechanism (Horbowicz et al., 2008). In freezing conditions, transparent co-pigment cooperates with anthocyanins, increasing their stability while intensifying their color. This process highly maintains and amplifies the anthocyanin content in fruits throughout storage. Indeed, 84% of the total antioxidant activity stemmed from anthocyanins, making them the main contributors (Skrovankova et al., 2015). Therefore, increased anthocyanin levels would raise the antioxidant activity and explain the results of this research.

On the contrary, some studies have shown that frozen conditions may either preserve or exert some adverse effects on fruit quality and functional properties. For example, anthocyanin levels remained statistically unchangeable in highbush blueberries (*Vaccinium corymbosum* L.) stored at -18 °C and -35 °C throughout the storage duration of six months (Scibisz and Mitek, 2007). Šamec et al. (2015) revealed different fluctuating outcomes of 10 small fruits stored at -20 °C for one year. The authors reported that many fruits maintained the TPC, TFC, and TAC after 12 months of frozen storage, whereas the hawthorn and strawberry fruits showed a decrease in most biochemical components (Šamec and Piljac-Žegarac, 2015).

Moreover, González et al. (2003) evaluated the effect of long-term frozen storage at -24 °C for 12 months on four different raspberry cultivars. During the first three months, the TPC in the 'Heritage' variety showed a significant decrease, while the 'Autumn Bliss' cultivar increased by 11% compared to fresh fruits. The TPC in the

'Zeva' and 'Rubi' varieties remained unchanged (González et al., 2003). Therefore, the diverse fluctuations in bioactive compounds greatly depend on the type of fruits, the cultivar, and the storage conditions, including the temperature and duration (González et al., 2003).

Likewise, other studies highlighted differences in outcomes compared to our results. Chaovanalikit and Wrolstad (2004) illustrated a significant reduction in the antioxidant activity of 'Bing' cherries stored at -23 °C after three months. However, a freezing temperature (-70 °C) enhanced the antioxidant potential by 14% (Chaovanalikit and Wrolstad, 2004). González et al. (2003) found a significant reduction in anthocyanins in four raspberries stored at -24 °C for three months (González et al. 2003). Vollmannová et al. (2009) also showed a decline in anthocyanins in most blueberry cultivars stored for three months at -18 °C (Vollmannová et al., 2009). In addition, Sahari et al. (2004) reported a similar drop in anthocyanins in strawberries stored at different temperatures (-12, -18, and -24 °C) for three months. The same study illustrated an increase in pH values, reaching 3.4 at -12 °C, potentially contributing to the observed decrease in anthocyanin content (Sahari et al., 2004), as suggested by various studies (Sablani, 2015; Wu et al., 2004). Also, Sistrunk et al. (1983) noted that anthocyanins degrade in fruits with a pH above 3.4 (Sistrunk et al., 1983). In contrast, the current research showed a decrease in pH levels by the end of the examined frozen storage, which might explain the increase in anthocyanin content. It could explain how frozen storage affects anthocyanin content in multiple ways, depending on the crop, cultivar, storage duration, temperature, and pH value (Sablani, 2015).

On the other hand, the increased titratable acidity content in our study may arise from the release of additional acids associated with cell wall softening and degradation (Hossain et al., 2014). In contrast, Sahari et al. (2004) showed that the acidity in strawberries varied during storage at different temperatures. At -12 °C, the acidity significantly declined over 90 days, reaching the lowest value (1.08). At -18 °C, the decline reached 1.15 after 60 days before remaining constant. However, a temperature of -24 °C retained the acidity stable after 30 days, and then a gradual decrease in a non-significant manner occurred (Sahari et al., 2004).

However, the decrease in total soluble solids could be due to a loss of water-soluble compounds due to the freezing and thawing process (Pistón et al., 2017). Similar findings by González et al. (2002) revealed a significant drop

of 4% in soluble solids in two different raspberry cultivars, 'Rubi' and 'Zeva,' stored at -24 °C for one year (González et al., 2002). In contrast, the same study revealed the opposite results in other cultivars, 'Heritage' and 'Autumn Bliss,' demonstrating a significant increase of 16% and 4%, respectively (González et al., 2002). Therefore, the total soluble solids strictly depended on the cultivar, environmental circumstances, and storage conditions (González et al., 2003). Similarly, the drop in weight at the end of the storage period can emanate from the loss in fruit water content, thus leading to an increase in the concentration of anthocyanins and phenols in blueberries (Ktenioudaki et al., 2021). The highest phenolic compounds in this research appeared in blueberries stored for three months but with the lowest fruit weight.

Conclusions

Blueberries at freezing temperatures (-20 °C) stored for three months resulted in a noteworthy enhancement of polyphenols, antioxidant activity, flavonoids, and anthocyanins content. This low-temperature storage maintained the titratable acidity, weight, and pH values for 60 days and more. By the end of the third month, a decrease occurred in the weight, pH, and total soluble solids accompanied by an increase in the titratable acidity. Therefore, we recommend using blueberries stored for more than two months in processing methods to fully exploit their richness in phenolic compounds and avoid issues related to their sour taste and reduced weight. In addition, further research can explore specific mechanisms underlying the changes in bioactive compounds during frozen storage, with potential health benefits and applications of frozen blueberries.

Conflict of Interest

The authors indicate no conflict of interest in this work.

References

- Adi DD, Oduro IN, Tortoe C. 2019. Physicochemical changes in plantain during normal storage ripening. *Scientific African* 6, 2468–2276. <https://doi.org/10.1016/j.sciaf.2019.e00164>
- Al Hasani S, Al-Attabi Z, Waly M, Al-Habsi N, Al-Subhi L, Shafiur Rahman M. 2023. Polyphenol and flavonoid stability of wild blueberry (*Sideroxylon mascatense*) during air- and freeze-drying and storage stability as a function of temperature. *Foods* 12, 871. <https://doi.org/10.3390/foods12040871>
- Bakowska-Barczak AM, Kolodziejczyk P. 2008. Evaluation of Saskatoon berry (*Amelanchier alnifolia*

Nutt.) cultivars for their polyphenol content, antioxidant properties, and storage stability. *Journal of Agricultural and Food Chemistry* 56, 9933–9940. <https://doi.org/10.1021/jf801887w>.

Brand-Williams W, Cuvelier ME, Berset C. 1995. Use of a free radical method to evaluate antioxidant activity. *LWT - Food Science and Technology* 28, 25–30. [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5)

Chaovanalikit A, Wrolstad RE. 2004. Total anthocyanins and total phenolics of fresh and processed cherries and their antioxidant properties. *Journal of Food Science* 69, FCT67–FCT72. <https://doi.org/10.1111/j.1365-2621.2004.tb17858.x>

Chen JY, Wen PF, Kong WF, Pan QH, Zhan JC, Li JM, Wan SB, Huang WD. 2006. Effect of salicylic acid on phenylpropanoids and phenylalanine ammonia-lyase in harvested grape berries. *Postharvest Biology and Technology* 40, 64–72. <https://doi.org/10.1016/j.postharvbio.2005.12.017>

De Ancos B, González EM, Cano MP. 2000. Ellagic acid, vitamin C, and total phenolic contents and radical scavenging capacity affected by freezing and frozen storage in raspberry fruit. *Journal of Agricultural and Food Chemistry* 48, 4565–4570. <https://doi.org/10.1021/jf0001684>

Duan Y, Tarafdar A, Chaurasia D, Singh A, Bhargava PC, Yang J, Li Z, Ni X, Tian Y, Li H, Awasthi MK. 2022. Blueberry fruit valorization and valuable constituents: a review. *International Journal of Food Microbiology* 381, 109890. <https://doi.org/10.1016/j.ijfoodmicro.2022.109890>

Edger PP, Iorizzo M, Bassil NV, Benevenuto J, Ferrão LFV, Giongo L, Hummer K, Lawas LMF, Leisner CP, Li C, Munoz PR, Ashrafi H, Atucha A, Babiker EM, Canales E, Chagné D, DeVetter L, Ehlenfeldt M, Espley RV, Gallardo K, Günther CS, Hardigan M, Hulse-Kemp AM, Jacobs M, Lila MA, Luby C, Main D, Mengist MF, Owens GL, Perkins-Veazie P, Polashock J, Pottorff M, Rowland LJ, Sims CA, Song G, Spencer J, Vorsa N, Yocca AE, Zalapa J. 2022. There and back again; historical perspective and future directions for *Vaccinium* breeding and research studies. *Horticulture Research* 9, uhac083. <https://doi.org/10.1093/hr/uhac083>

Giusti MM, Wrolstad RE. 2003. Acylated anthocyanins from edible sources and their applications in food systems. *Biochemical Engineering Journal* 14, 217–225. [https://doi.org/10.1016/S1369-703X\(02\)00221-8](https://doi.org/10.1016/S1369-703X(02)00221-8)

González E, De Ancos B, Cano P. 2002. Preservation of raspberry fruits by freezing: physical, physico-chemical and sensory aspects. *European Food Research and Technology* 215, 497–503. <https://doi.org/10.1007/s00217-002-0600-4>

González EM, De Ancos B, Cano MP. 2003. Relation between bioactive compounds and free radical-scavenging capacity in berry fruits during frozen storage: bioactive compounds in frozen-stored berry fruits. *Journal of the Science of Food and Agriculture* 83, 722–726. <https://doi.org/10.1002/jsfa.1359>

- Horbowicz M, Kosson R, Grzesiuk A, Dębski H. 2008. Anthocyanins of fruits and vegetables - their occurrence, analysis and role in human nutrition. *Journal of Fruit and Ornamental Plant Research* 68, 5-22. <https://doi.org/10.2478/v10032-008-0001-8>
- Hossain MdA, Rana MdM, Kimura Y, Roslan HA. 2014. Changes in biochemical characteristics and activities of ripening associated enzymes in mango fruit during the storage at different temperatures. *BioMed Research International* 232969. <https://doi.org/10.1155/2014/232969>
- King ES, Noll A, Glenn S, Bolling BW. 2022. Refrigerated and frozen storage impact aronia berry quality. *Food Production, Processing and Nutrition* 4, 3. <https://doi.org/10.1186/s43014-021-00080-y>
- Ktenioudaki A, O'Donnell CP, Emond JP, do Nascimento Nunes MC. 2021. Blueberry supply chain: critical steps impacting fruit quality and application of a boosted regression tree model to predict weight loss. *Postharvest Biology and Technology* 179, 111590. <https://doi.org/10.1016/j.postharvbio.2021.111590>
- Lamaison JL, Carnat A. 1990. Teneur en principaux flavonoïdes des fleurs et des feuilles de *Crataegus monogyna* Jacq. et de *Crataegus laevigata* (Poiret) DC. (Rosaceae). *Pharmaceutica Acta Helvetiae* 65, 315-320.
- Mahajan PV, Caleb OJ, Singh Z, Watkins CB, Geyer M. 2014. Postharvest treatments of fresh produce. *Philosophical Transactions of the Royal Society* 372, 20130309. <https://doi.org/10.1098/rsta.2013.0309>
- Matiacevich S, Celis Cofré D, Silva P, Enrione J, Osorio F. 2013. Quality parameters of six cultivars of blueberry using computer vision. *International Journal of Food Science* 1-8. <https://doi.org/10.1155/2013/419535>
- Mensor LL, Menezes FS, Leitão GG, Reis AS, Santos TCD, Coube CS, Leitão SG. 2001. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method: antioxidant activity in Brazilian plants. *Phytotherapy Research* 15, 127-130. <https://doi.org/10.1002/ptr.687>
- Nielsen SS. 2017. *Food Analysis Laboratory Manual*, (3rd ed.), Food Science Text Series. Springer International Publishing, Cham. <https://doi.org/10.1007/978-3-319-44127-6>
- Olivas-Aguirre F, Rodrigo-García J, Martínez-Ruiz N, Cárdenas-Robles A, Mendoza-Díaz S, Álvarez-Parrilla E, González-Aguilar G, De La Rosa L, Ramos-Jiménez A, Wall-Medrano A. 2016. Cyanidin-3-O-glucoside: physical-chemistry, foodomics and health effects. *Molecules* 21, 1264. <https://doi.org/10.3390/molecules21091264>
- Piljac-Žegarac J, Šamec D. 2011. Antioxidant stability of small fruits in postharvest storage at room and refrigerator temperatures. *Food Research International* 44, 345-350. <https://doi.org/10.1016/j.foodres.2010.09.039>
- Pistón F, Pérez AG, Sanz C, Refoyo A. 2017. Relationship between sugar content and °Brix as influenced by cultivar and ripening stages of strawberry. *Acta Horticulturae* 491-496. <https://doi.org/10.17660/ActaHortic.2017.1156.72>
- Reque P.M, Steffens R.S, Jablonski A, Flôres S.H, Rios A.D.O, De Jong E.V. 2014. Cold storage of blueberry (*Vaccinium* spp.) fruits and juice: anthocyanin stability and antioxidant activity. *Journal of Food Composition and Analysis* 33, 111-116. <https://doi.org/10.1016/j.jfca.2013.11.007>
- Rodriguez-Mateos A, Cifuentes-Gomez T, Tabatabaee S, Lecras C, Spencer JPE. 2012. Procyanidin, anthocyanin, and chlorogenic acid contents of highbush and lowbush blueberries. *Journal of Agricultural and Food Chemistry* 60, 5772-5778. <https://doi.org/10.1021/jf203812w>
- Sablani S.S. 2015. "Freezing of Fruits and Impact on Anthocyanins" in *Processing and Impact on Active Components in Food*. Elsevier, 147-156. <https://doi.org/10.1016/B978-0-12-404699-3.00018-4>
- Sahari MA, Mohsen Boostani F, Zohreh Hamidi E. 2004. Effect of low temperature on the ascorbic acid content and quality characteristics of frozen strawberry. *Food Chemistry* 86, 357-363. <https://doi.org/10.1016/j.foodchem.2003.09.008>
- Šamec D, Piljac-Žegarac J. 2015. Fluctuations in the levels of antioxidant compounds and antioxidant capacity of ten small fruits during one year of frozen storage. *International Journal of Food Properties* 18, 21-32. <https://doi.org/10.1080/10942912.2013.790423>
- Sánchez-Rangel JC, Benavides J, Heredia JB, Cisneros-Zevallos L, Jacobo-Velázquez DA. 2013. The Folin-Ciocalteu assay revisited: improvement of its specificity for total phenolic content determination. *Analytical Methods* 5, 5990. <https://doi.org/10.1039/c3ay41125g>
- Scibisz I, Mitek M. 2007. The changes of antioxidant properties in highbush blueberries (*Vaccinium corymbosum* L.) during freezing and long-term frozen storage. *Acta Scientiarum Polonorum: Technologia Alimentaria* 6, 75-82.
- Seeram NP. 2008. Berry fruits: compositional elements, biochemical activities, and the impact of their intake on human health, performance, and disease. *Journal of Agricultural and Food Chemistry* 56, 627-629. <https://doi.org/10.1021/jf071988k>
- Silva S, Costa EM, Veiga M, Morais RM, Calhau C, Pintado M. 2020. Health promoting properties of blueberries: a review. *Critical Reviews in Food Science and Nutrition* 60, 181-200. <https://doi.org/10.1080/10408398.2018.1518895>
- Sistrunk WA, Wang RC, Morris JR. 1983. Effect of combining mechanically harvested green and ripe puree and sliced fruit, processing methodology and frozen storage on quality of strawberries. *Journal of Food Science* 48, 1609-1612. <https://doi.org/10.1111/j.1365-2621.1983.tb05043.x>
- Skrovankova S, Sumczynski D, Mlcek J, Jurikova T,

Sochor J. 2015. Bioactive compounds and antioxidant activity in different types of berries. *International Journal of Molecular Sciences* 16, 24673–24706. <https://doi.org/10.3390/ijms161024673>

Smith MAL, Marley KA, Seigler D, Singletary KW, Meline B. 2000. Bioactive properties of wild blueberry fruits. *Journal of Food Science* 65, 352–356. <https://doi.org/10.1111/j.1365-2621.2000.tb16006.x>

Tietel Z, Lewinsohn E, Fallik E, Porat R. 2012. Importance of storage temperatures in maintaining flavor and quality of mandarins. *Postharvest Biology and Technology* 64, 175–182. <https://doi.org/10.1016/j.postharvbio.2011.07.009>

Veberic R, Slatnar A, Bizjak J, Stampar F, Mikulic-Petkovsek M. 2015. Anthocyanin composition of different wild and cultivated berry species. *LWT - Food Science and Technology* 60, 509–517. <https://doi.org/10.1016/j.lwt.2014.08.033>

Vollmannová A, Tóth T, Urminská D, Poláková Z, Timoracká M, Margitanová E. 2009. Anthocyanins content in blueberries (*Vaccinium corymbosum* L.) in relation to freezing duration. *Czech Journal of Food Sciences* 27, S204–S206. <https://doi.org/10.17221/1069-CJFS>

Vrhovsek U, Masuero D, Palmieri L, Mattivi F. 2012. Identification and quantification of flavonol glycosides in cultivated blueberry cultivars. *Journal of Food Composition and Analysis* 25, 9–16. <https://doi.org/10.1016/j.jfca.2011.04.015>

Wu X, Gu L, Prior RL, McKay S. 2004. Characterization of anthocyanins and proanthocyanidins in some cultivars of *Ribes*, *Aronia*, and *Sambucus* and their antioxidant capacity. *Journal of Agricultural and Food Chemistry* 52, 7846–7856. <https://doi.org/10.1021/jf0486850>

Young JC, Abdel-Aal ESM. 2009. "Anthocyanins" in *Healthgrain Methods*. Elsevier, 141–165. <https://doi.org/10.1016/B978-1-891127-70-0.50015-7>