



## Enhancing Morpho-phytochemical Properties of *Catharanthus roseus* L. var. 'Ocellatus' via Plant Growth Regulators

Elham Jafarpour, Azizollah Kheiry\*, Mohsen SaniKhani, Farhang Razavi

1 Department of Horticultural Science, Faculty of Agriculture, University of Zanjan, Zanjan, Iran

### ARTICLE INFO

#### Article history:

Received: 26 November 2023,  
Received in revised form: 1 April 2024,  
Accepted: 3 April 2024

#### Article type:

Research paper

#### Keywords:

Antioxidant capacity,  
Auxins,  
*Catharanthus roseus* L var. Ocellata,  
Medicinal plant,  
Tissue culture

### COPYRIGHT

© 2023 The author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other medium is permitted, provided the original author(s) and source are cited, in accordance with accepted academic practice. No permission is required from the authors or the publishers.

### ABSTRACT

*Catharanthus roseus* L. is a valuable medicinal plant renowned for its potent anticancer compounds, i.e., vincristine and vinblastine. The efficient mass production of this plant can be challenging due to limitations in conventional propagation methods. Tissue culture techniques offer a promising alternative, and optimizing plant growth regulator (PGR) treatments can play a crucial role in achieving successful *in vitro* propagation. In this study, we investigated the effects of several PGR combinations, i.e., NAA, BAP, and 2,4-D, on the morpho-phytochemical attributes and antioxidant activity of *in vitro*-grown *Catharanthus roseus* L. var. 'Ocellata'. The plants were arranged in a completely randomized design with three replications. The results demonstrated that applying PGRs, either alone or in combination, significantly improved the morphological characteristics. The NAA treatment caused the highest plant height, number of branches and leaves, and root diameter. Notably, the treatment with 2,4-D resulted in the highest photosynthetic pigment content compared to the control. Moreover, the treatments with NAA + BAP and NAA + BAP + 2,4-D exhibited the highest levels of total phenolic compounds (TPC) and total flavonoid content (TFC) in both roots and leaves. Regarding antioxidant activity, the DPPH radical scavenging assay revealed the highest radical scavenging percentages, i.e., 74.21% (NAA + 2,4-D) in leaf samples and 78.08% (NAA + 2,4-D) in root samples. These findings emphasize the potency of PGRs in optimizing tissue culture protocols for *Catharanthus roseus* L., thus facilitating the production of superior-quality plants with enriched medicinal properties. This study contributes to the advancement of sustainable and efficient cultivation of bioactive compounds from *Catharanthus roseus* L. using tissue culture techniques.

### Introduction

*Catharanthus roseus* L., commonly known as Madagascar periwinkle, is a medicinal plant renowned for its production of valuable anticancer compounds, vincristine, and vinblastine (Kabesh et al., 2015; Pereira et al., 2009). These compounds possess significance in treating various types of

cancer, making *Catharanthus roseus* a highly sought-after plant in the pharmaceutical industry (Chen et al., 2018; Yamamura et al., 2017; Montagna et al., 2017; Gajalakshmi et al., 2013; Marcone et al., 1997; Pandey-Rai et al., 2006). However, conventional methods of propagation, such as sexual reproduction and cutting, exhibit low success

\*Corresponding author's email: [kheiry@znu.ac.ir](mailto:kheiry@znu.ac.ir)

rates and pose challenges for large-scale production (Bakrudeen et al., 2011). Tissue culture techniques have partly solved these problems and have emerged as a promising approach for the efficient and sustainable production of this valuable plant species (Sandhya et al., 2016).

Tissue culture, conducted under controlled laboratory conditions, provides an ideal environment conducive to propagating plant material rapidly and accurately while maintaining genetic uniformity and desired traits (Anand, 2010; Smetanska, 2008). In tissue culture, optimizing culture media components, including plant growth regulators (PGRs), can vitally assist in enhancing growth, development, and secondary metabolite production under *in vitro*-grown plants (Roberts, 2012).

PGRs, such as auxins and cytokinin, and their combinations, reportedly regulate various physiological processes in plants (Yong et al., 2009). They influence cell division, differentiation, and organogenesis, thus affecting the growth and development of tissue-cultured plants (Mitra et al., 2016). Moreover, PGRs have reportedly amended the biosynthesis of secondary metabolites, including phenolic components, flavonoids, and alkaloids, which are responsible for the therapeutic properties of medicinal plants (Derbassi et al., 2022; Ochatt et al., 2022).

In this study, our objective was to investigate the effects of different PGR treatments on the morphological and phytochemical characteristics of *in vitro*-grown *Catharanthus roseus* L. var. 'Ocellata' plants. We sought to optimize the tissue culture conditions to increase the production of valuable bioactive compounds with potential therapeutic applications. By evaluating various combined and separate PGR treatments, including NAA (Naphthalene acetic acid), BAP (6-benzylaminopurine), and 2,4-D (2,4-dichlorophenoxyacetic acid), we aimed to elucidate the specific roles of these regulators in enhancing plant growth, biomass accumulation, and secondary metabolite synthesis.

## Materials and Methods

### **Plant materials and explant preparation**

Nodal shoot explants (with two pairs of leaves) were obtained from healthy *Catharanthus roseus* L. plants grown in the research greenhouse at the University of Zanjan. The explants were immediately transferred to the Horticulture Biotechnology and Tissue Culture Laboratory for further processing.

### **Sterilization protocol**

The explants were subjected to a sterilization protocol to eliminate any surface contaminants. The protocol involved washing the explants with running tap water for 5 min, followed by immersion in a solution of 70% ethanol for 1 min, and subsequently in 5% sodium hypochlorite for 10 min. The explants were then rinsed three times with sterile distilled water (Hernández-Domínguez et al., 2004).

### **Culture medium and treatment setup**

After sterilization, the explants were placed on a culture medium based on Murashige and Skoog (MS) (Classic Murashige and Skoog, 1962) medium supplemented with different PGRs as treatments. The PGR treatments were NAA (1 mg L<sup>-1</sup>), (1 mg L<sup>-1</sup> + BAP (1 mg L<sup>-1</sup>), NAA (1 mg L<sup>-1</sup> + 2,4-D (0.5 mg L<sup>-1</sup>), NAA (1 mg L<sup>-1</sup> + BAP (1 mg L<sup>-1</sup> + 2,4-D (0.5 mg L<sup>-1</sup>), BAP (1 mg L<sup>-1</sup>), BAP (1 mg L<sup>-1</sup> + 2,4-D (0.5 mg L<sup>-1</sup>), 2,4-D (0.5 mg L<sup>-1</sup>). A control treatment without any PGRs was also included. The research was conducted as a completely randomized design (CRD) with three replications.

### **Incubation and growth conditions**

Following the treatment setup, the explants were incubated in a growth chamber at a temperature of 25 °C under a 16 h photoperiod provided by white fluorescent lights with a photon flux density of 70 μmol m<sup>-2</sup> s<sup>-1</sup>.

### **Measurement of morphological traits**

After 90 days of growth in the tissue culture environment, various morphological traits were measured, including plant height, stem diameter, intermediate length, root diameter, root length, number of branches (NB), number of leaves, fresh yield, dry yield, tissue water content (shoot and root water content), tissues dry matter content (shoot and root dry matter content), shoot fresh yield, root fresh yield, shoot dry yield, and root dry yield.

### **Measurement of photosynthetic pigments**

Chlorophyll a, chlorophyll b, and carotenoid contents were evaluated according to a method described by Arnone (1949).

### **Measurement of phytochemicals, determination of total phenolic and flavonoid content**

Phytochemical attributes such as total phenolic content (TPC) and total flavonoid content (TFC) were assayed. Methanolic extracts of shoot and root tissues were prepared and the TPC was evaluated

using the Folin-Ciocalteu reagent method and calibrated against Gallic acid standards, following a relevant protocol (Singleton and Rossi, 1965). The results were expressed as mg of Gallic acid equivalents (GAE mg g<sup>-1</sup>) per g of dry weight extract. A spectrophotometer measured the absorbance at 765 nm.

The spectrophotometric colorimetric method (Kähkönen et al., 1999) determined the TFC. The reaction mixture was prepared in a test tube by adding 500 µL of the extract solution, 100 µL of 10% aluminum chloride solution, 100 µL of 1 M potassium acetate solution, and 2.8 mL of distilled water. The samples were incubated for 40 min at room temperature, and the absorbance was measured at 415 nm using a spectrophotometer (T-60, PG Instrument UK). Quercetin was used to plot the standard curve, and the results were expressed as milligrams of quercetin equivalents (QE mg g<sup>-1</sup>) per gram of extract.

#### **Antioxidant capacity determination**

The antioxidant capacity of the plant extracts was assessed using three different methods: DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging assay, reducing power assay, and phosphomolybdenum assay (TAC assay).

#### **DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging assay**

The DPPH radical scavenging assay was performed according to a relevant method (Hatano et al., 1988). A stock solution of DPPH was prepared by dissolving 4 mg in 100 mL of methanol and kept at 20 °C for further use. A mixture of 250 µL of plant samples and 750 µL of DPPH aliquot was incubated for 15 min at room temperature in the dark. Optical density was measured at a wavelength of 517 nm. The antioxidant capacity was calculated using the following equation:

$$\text{Free radical scavenging activity (\%)} = \frac{\text{Control absorbance} - \text{Sample absorbance}}{\text{Control absorbance}} \times 100$$

#### **Reducing power assay**

Reducing power activity was determined using a method described in the literature (Oyaizu, 1986). A mixture of 0.5 mL plant extract, 0.5 mL 0.2 M phosphate buffer (pH 6.6), and 0.5 mL potassium ferricyanide (10 mg L<sup>-1</sup>) was incubated at 50 °C for 20 min. Then, 0.5 mL of trichloroacetic acid (TCA) (100 mg L<sup>-1</sup>) was added to the solution, and the mixture

was centrifuged at 3000 rpm for 10 min. Then, 0.5 mL of the supernatant was diluted with 0.5 mL of pure water and mixed with 0.1 mL of FeCl<sub>3</sub> (0.1%). The optical density was measured at 700 nm after 10 min. Methanol (80%) was used as a blank, and a solution mixture without plant extract served as the control under the same preparation conditions for each plant sample.

#### **Phosphomolybdenum assay or total antioxidant capacity (TAC) assay**

Total antioxidant capacity of plant samples was determined using the phosphomolybdenum assay as described in the literature (Prieto et al., 1999). The phosphomolybdenum reagent solution was prepared by mixing 1 mL of Na<sub>3</sub>PO<sub>4</sub> (28 mM), H<sub>2</sub>SO<sub>4</sub> (0.6 M), and ammonium molybdate (4 mM) with 0.3 mL of the plant extract. The reaction mixture was heated at 95 °C in a water bath for 90 min, covered with silver foil to avoid direct light exposure. After cooling them down to room temperature, the absorbance was measured at 695 nm using a spectrophotometer. All treatments occurred in triplicates.

$$\% \text{ TAC inhibition} = \frac{\text{Abs sample} - \text{Abs control}}{\text{Abs sample}} \times 100$$

#### **Statistical analysis**

All data were subjected to one-way analysis of variance (ANOVA) using SAS statistical analysis software version 9.4. Data were expressed as mean values ± standard deviation (SD), and significant differences were determined among the treatment groups using LSD tests (P≤0.05).

## **Results**

### **Morphological attributes**


Full plantlet regeneration of the typical shoot explants on MS medium supplemented with various PGRs occurred in the third month of in vitro culture (Figs. 1 and 2).

Compared to the control, all measured morphological traits showed significant differences in all treatments. Regarding most of the desired traits, the PGRS treatments improved morphological attributes (Table 1). The highest mean values of height (71.75 mm), number of lateral branches (5), number of leaves (40), and internode length (9.71 mm) occurred in response to the 1 mg L<sup>-1</sup> NAA treatment. Regarding plant diameter, root diameter, and length, maximum values (4.67 mm, 3.67 mm, and 115.97 mm) occurred in response to the 1 mg L<sup>-1</sup>




$1 \text{ mg L}^{-1}$  BAP +  $0.5 \text{ mg L}^{-1}$  2,4-D,  $1 \text{ mg L}^{-1}$  NAA +  $1 \text{ mg L}^{-1}$  BAP, and  $0.5 \text{ mg L}^{-1}$  2,4-D treatments, respectively. Meanwhile, the smallest root diameter and length ( $0.53 \text{ mm}$  and  $10.66 \text{ mm}$ ) were in the control group. The fresh and dry yield and its components, such as fresh and dry yield of the roots and shoots, had the

highest significant values ( $8662.43 \text{ mg}$ ,  $709.7 \text{ mg}$ ,  $3051.66 \text{ mg}$ ,  $5610.77 \text{ mg}$ ,  $236.5 \text{ mg}$ , and  $473.2 \text{ mg}$ ) in MS medium supplemented with  $1 \text{ mg L}^{-1}$  NAA +  $1 \text{ mg L}^{-1}$  BAP (Table 1).



**Fig. 1.** Shoot regeneration of the typical shoot explant of *C. roseus* L. var. 'Ocellatus' on MS medium supplemented with different PGRs (a: NAA, b: NAA + BAP, c: NAA + 2,4-D, d: NAA + BAP + 2,4-D, e: BAP, f: BAP + 2,4-D, g: 2,4-D, h: control). NAA: 1-Naphthaleneacetic acid, BAP: 6-Benzylaminopurine, 2,4-D: 2,4-Dichlorophenoxyacetic acid.



**Fig. 2.** Root system regeneration of the typical shoot explant of *C. roseus* L. var. 'Ocellatus' on MS medium supplemented with different PGRs (a: NAA, b: NAA + BAP, c: NAA + 2,4-D, d: NAA + BAP + 2,4-D, e: BAP, f: BAP + 2,4-D, g: 2,4-D, h: control). NAA: 1-Naphthaleneacetic acid, BAP: 6-Benzylaminopurine, 2,4-D: 2,4-Dichlorophenoxyacetic acid.

**Table 1.** Morphological traits of *C. roseus* var. 'Ocellatus' on MS medium supplemented with different PGRs.

Treatments	Height (mm)	NB	Plant Diameter (mm)	NL	IL (mm)	RD (mm)	RL (mm)
1 mg L <sup>-1</sup> NAA	71.75 ± 0.09 <sup>a</sup>	5 ± 0 <sup>a</sup>	2.75 ± 0.01 <sup>c</sup>	40 ± 1 <sup>a</sup>	9.71 ± 0.29 <sup>a</sup>	3.04 ± 0.58 <sup>a</sup>	22.76 ± 0.29 <sup>d</sup>
1 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP	68.96 ± 18.5 <sup>ab</sup>	3 ± 1 <sup>b</sup>	3.29 ± 0.9 <sup>bc</sup>	29 ± 8.7 <sup>b</sup>	7.55 ± 1.21 <sup>abc</sup>	3.67 ± 0.59 <sup>a</sup>	70.14 ± 9.5 <sup>b</sup>
1 mg L <sup>-1</sup> NAA + 0.5 mg L <sup>-1</sup> 2,4-D	57.06 ± 0.6 <sup>bcd</sup>	1.66 ± 0.57 <sup>cd</sup>	2.71 ± 0.04 <sup>c</sup>	25 ± 1 <sup>b</sup>	7.72 ± 0.05 <sup>ab</sup>	0.46 ± 0.05 <sup>b</sup>	24 ± 4 <sup>d</sup>
1 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP + 0.5 mg L <sup>-1</sup> 2,4-D	41.02 ± 1.5 <sup>e</sup>	2 ± 0 <sup>c</sup>	3.5 ± 0.68 <sup>bc</sup>	22.33 ± 5.03 <sup>b</sup>	5.49 ± 0.96 <sup>cd</sup>	3.01 ± 1.9 <sup>a</sup>	24.17 ± 3.7 <sup>cd</sup>
1 mg L <sup>-1</sup> BAP	62.18 ± 5.05 <sup>abc</sup>	3 ± 0 <sup>b</sup>	2.92 ± 0.19 <sup>c</sup>	28.33 ± 1.5 <sup>b</sup>	7.48 ± 2.08 <sup>bc</sup>	0.74 ± 0.12 <sup>b</sup>	45.28 ± 26.56 <sup>c</sup>
1 mg L <sup>-1</sup> BAP + 0.5 mg L <sup>-1</sup> 2,4-D	45.01 ± 2.8 <sup>de</sup>	1 ± 0 <sup>d</sup>	4.67 ± 0.4 <sup>a</sup>	23 ± 3 <sup>b</sup>	6.08 ± 0.82 <sup>bcd</sup>	1.44 ± 0.43 <sup>b</sup>	8.50 ± 1.2 <sup>d</sup>
0.5 mg L <sup>-1</sup> 2,4-D	51.61 ± 5.7 <sup>cde</sup>	1.33 ± 0.57 <sup>cd</sup>	3.21 ± 0.05 <sup>bc</sup>	24.66 ± 5.03 <sup>b</sup>	5.17 ± 2.06 <sup>d</sup>	0.76 ± 0.09 <sup>b</sup>	115.97 ± 18.1 <sup>a</sup>
Control	24.49 ± 3.1 <sup>f</sup>	1 ± 0 <sup>d</sup>	3.89 ± 0.41 <sup>ab</sup>	22.66 ± 2.3 <sup>b</sup>	2.29 ± 0.99 <sup>c</sup>	0.53 ± 0.03 <sup>b</sup>	10.66 ± 7.17 <sup>d</sup>
LSD	12.57	0.79	0.79	7.37	2.18	1.34	21.26
Significant Treatment	df						
	7	**	**	**	**	**	**
CV		13.76	20.28	13.58	15.85	19.63	15.50

Data was expressed as means ± standard deviation (n = 3). Different superscript letters in the same column indicate significant differences according to LSD ( $p \leq 0.05$ ), \*\* and \* are significant at the 0.01 and 0.05 levels. (Nb: number of branches, NL: number of leaves, IL: intermediate length, RD: root diameter, RL: root length). Data are mean values of n = 3. Mean values followed by different letters indicate they are significantly different by the LSD test ( $p < 0.05$ ). NAA: 1-Naphthaleneacetic acid, BAP: 6-Benzylaminopurine, 2,4-D: 2,4-Dichlorophenoxyacetic acid.

Table 1. (Continued)

Treatments	FY (mg)	DY (mg)	SHTW (%)	RTW (%)	DRM (%)	DSHM (%)	SHFY (mg)	RFY (mg)	SHDY (mg)	RDY (mg)
1 mg L <sup>-1</sup> NAA	7021.9 ± 7.8 <sup>c</sup>	411.23 ± 0.45 <sup>c</sup>	94.42 ± 2.25 <sup>a</sup>	93.98 ± 0.08 <sup>a</sup>	6.04 ± 0.08 <sup>c</sup>	5.58 ± 2.25 <sup>h</sup>	2550.16 ± 5.15 <sup>c</sup>	4471.7 ± 5.8 <sup>c</sup>	142.26 ± 0.3 <sup>d</sup>	268.96 ± 0.15 <sup>c</sup>
1 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP	8662.43 ± 8.6 <sup>a</sup>	709.7 ± 1.1 <sup>a</sup>	92.25 ± 1.31 <sup>e</sup>	91.56 ± 1.56 <sup>e</sup>	8.43 ± 1.56 <sup>c</sup>	7.75 ± 1.31 <sup>d</sup>	3051.66 ± 10.02 <sup>a</sup>	5610.77 ± 17.9 <sup>a</sup>	236.5 ± 0.8 <sup>a</sup>	473.2 ± 1.6 <sup>a</sup>
1 mg L <sup>-1</sup> NAA + 0.5 mg L <sup>-1</sup> 2,4-D	8028.2 ± 9.1 <sup>b</sup>	527.66 ± 1.2 <sup>b</sup>	93.43 ± 0 <sup>b</sup>	93.42 ± 0 <sup>b</sup>	6.57 ± 0 <sup>d</sup>	6.57 ± 0 <sup>g</sup>	2930 ± 10 <sup>b</sup>	5098.2 ± 2 <sup>b</sup>	192.4 ± 1.3 <sup>b</sup>	335.2 ± 0.17 <sup>b</sup>
1 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP + 0.5 mg L <sup>-1</sup> 2,4-D	4993 ± 28.7 <sup>d</sup>	318.7 ± 2.6 <sup>d</sup>	92.43 ± 0.54 <sup>d</sup>	94.17 ± 0.71 <sup>a</sup>	5.82 ± 0.71 <sup>e</sup>	7.56 ± 0.54 <sup>e</sup>	1603.1 ± 6.5 <sup>e</sup>	3389.9 ± 27.4 <sup>d</sup>	121.2 ± 0.5 <sup>e</sup>	197.5 ± 2.3 <sup>d</sup>
1 mg L <sup>-1</sup> BAP	1602.97 ± 14.27 <sup>g</sup>	132.86 ± 1.4 <sup>g</sup>	1.43 ± 1.62 <sup>f</sup>	94 ± 2.47 <sup>a</sup>	5.99 ± 2.47 <sup>c</sup>	8.56 ± 1.62 <sup>c</sup>	1429.6 ± 10.5 <sup>f</sup>	173.37 ± 4.08 <sup>g</sup>	122.4 ± 1.2 <sup>e</sup>	10.39 ± 0.35 <sup>g</sup>
1 mg L <sup>-1</sup> BAP + 0.5 mg L <sup>-1</sup> 2,4-D	3038.30 ± 48.37 <sup>c</sup>	302.83 ± 4.8 <sup>c</sup>	90.93 ± 0.78 <sup>h</sup>	88.75 ± 1.2 <sup>d</sup>	11.24 ± 1.2 <sup>b</sup>	9.07 ± 0.78 <sup>a</sup>	1783.76 ± 11.3 <sup>d</sup>	1254.5 ± 37.10 <sup>e</sup>	161.76 ± 0.9 <sup>c</sup>	141.06 ± 3.9 <sup>e</sup>
0.5 mg L <sup>-1</sup> 2,4-D	1970.30 ± 11.7 <sup>f</sup>	171.7 ± 1.5 <sup>f</sup>	91.24 ± 2.07 <sup>g</sup>	91.47 ± 2.13 <sup>c</sup>	8.52 ± 2.13 <sup>c</sup>	8.75 ± 2.07 <sup>b</sup>	1606.46 ± 10.1 <sup>e</sup>	363.8 ± 5.3 <sup>f</sup>	140.6 ± 1.04 <sup>d</sup>	31.03 ± 1.16 <sup>f</sup>
Control	864.53 ± 16.45 <sup>h</sup>	64.76 ± 0.6 <sup>h</sup>	93.11 ± 1.67 <sup>c</sup>	86.31 ± 5.18 <sup>e</sup>	13.68 ± 5.18 <sup>a</sup>	6.88 ± 1.67 <sup>f</sup>	787.53 ± 15.2 <sup>g</sup>	77 ± 1.5 <sup>h</sup>	54.23 ± 0.76 <sup>f</sup>	10.53 ± 0.40 <sup>g</sup>
LSD	38.7	3.81	0.04	0.35	0.35	0.04	17.8	30.83	1.6	3.08
Significant Treatment	df	**	**	**	**	**	**	**	**	**
CV	7	0.49	0.66	0.02	0.22	2.49	0.35	0.52	0.69	0.97

Data was expressed as means ± standard deviation (n = 3). Different superscript letters in the same column indicate significant differences according to LSD ( $p \leq 0.05$ ), \*\* and \* are significant at the 0.01 and 0.05 levels. (FY: fresh yield, DY: dry yield, SHTW: shoot tissue water content, RTW: root tissue water content, DRM: dry root matter, DSHM: dry shoot matter, SHFY: shoot fresh yield, RFY: root fresh yield, SHDY: shoot dry yield, RDY: root dry yield. Data are mean values of n = 3. Mean values followed by different letters indicate they are significantly different by the LSD test ( $p < 0.05$ ). NAA: 1-Naphthaleneacetic acid, BAP: 6-Benzylaminopurine, 2,4-D: 2,4-Dichlorophenoxyacetic acid.

**Photosynthetic pigments**

Data analysis indicated that MS medium supplemented with 0.5 mg L<sup>-1</sup> 2,4-D produced the highest chlorophyll a, chlorophyll b, and carotenoid

content (2.62, 0.63, 0.36 mg g<sup>-1</sup> FW). The lowest mean values for these variables (1.18, 0.26, 0.20 mg g<sup>-1</sup> FW, respectively) occurred in the control (Table 2).

**Table 2.** Photosynthetic pigments content of *C. roseus* L. var. 'Ocellatus' on MS medium supplemented with different PGRs.

Treatments	Chlorophyll a (mg g <sup>-1</sup> FW)	Chlorophyll b (mg g <sup>-1</sup> FW)	Carotenoid (mg g <sup>-1</sup> FW)
1 mg L <sup>-1</sup> NAA	2.05 ± 0.25 <sup>ab</sup>	0.48 ± 0.11 <sup>ab</sup>	0.32 ± 0.04 <sup>ab</sup>
1 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP	1.97 ± 0.64 <sup>ac</sup>	0.43 ± 0.14 <sup>ac</sup>	0.29 ± 0.07 <sup>ac</sup>
1 mg L <sup>-1</sup> NAA + 0.5 mg L <sup>-1</sup> 2,4-D	1.47 ± 0.11 <sup>bc</sup>	0.36 ± 0.02 <sup>bc</sup>	0.22 ± 0.01 <sup>bc</sup>
1 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP + 0.5 mg L <sup>-1</sup> 2,4-D	1.78 ± 0.75 <sup>ac</sup>	0.36 ± 0.13 <sup>bc</sup>	0.27 ± 0.1 <sup>ac</sup>
1 mg L <sup>-1</sup> BAP	1.99 ± 0.57 <sup>ac</sup>	0.41 ± 0.12 <sup>bc</sup>	0.29 ± 0.06 <sup>ac</sup>
1 mg L <sup>-1</sup> BAP + 0.5 mg L <sup>-1</sup> 2,4-D	1.81 ± 0.44 <sup>ac</sup>	0.35 ± 0.07 <sup>bc</sup>	0.27 ± 0.05 <sup>ac</sup>
0.5 mg L <sup>-1</sup> 2,4-D	2.6 ± 0.42 <sup>a</sup>	0.63 ± 0.16 <sup>a</sup>	0.36 ± 0.06 <sup>a</sup>
Control	1.18 ± 0.39 <sup>c</sup>	0.26 ± 0.08 <sup>c</sup>	0.2 ± 0.07 <sup>c</sup>
LSD	0.8501	0.2007	0.1153
Significant	df		
Treatment	7	*	*
CV		26.34	27.91
			23.68

Data are mean values of n = 3. Mean values followed by different letters indicate they are significantly different by the LSD test ( $p < 0.05$ ). NAA: 1-Naphthaleneacetic acid, BAP: 6-Benzylaminopurine, 2,4-D: 2,4-Dichlorophenoxyacetic acid.

**Phytochemical attributes**


TFC varied among the treatment groups and had significant differences (Table 3), ranging from 128.82 (1 mg L<sup>-1</sup> BAP + 0.5 mg L<sup>-1</sup> 2,4-D) to 36.95 (1 mg L<sup>-1</sup> NAA + 0.5 mg L<sup>-1</sup> 2,4-D) in leaf samples and from

24.50 (1 mg L<sup>-1</sup> NAA + 1 mg L<sup>-1</sup> BAP + 0.5 mg L<sup>-1</sup> 2,4-D) to 0.7 (control) in root samples (Fig. 3A). Maximum TPC in the methanolic extract of the leaf and root samples (respectively, 394.5 and 306.17) occurred in response to 1 mg L<sup>-1</sup> NAA + 1 mg L<sup>-1</sup> BAP treatment (Fig. 3B).

**Table 3.** Phytochemical traits of *C. roseus* L. var. 'Ocellatus' on MS medium supplemented with different PGRs.

SOV	DF	Mean Square (MS)									
		TFL	TFR	TPL	TPR	DPPHL	DPPHR	TACL	TACR	PFRAP <sub>L</sub>	PFRAPR
Treatment	7	2913.20 <sup>**</sup>	216.71 <sup>**</sup>	10559.42 <sup>**</sup>	23363.27 <sup>**</sup>	26.35 <sup>*</sup>	600.06 <sup>**</sup>	67.39 <sup>**</sup>	297.70 <sup>**</sup>	122.67 <sup>**</sup>	422.96 <sup>**</sup>
Error	16	21.03	1.42	24.10	1.42	10.14	2.34	2.86	2.34	7.87	7.04
LSD		7.93	2.06	8.49	2.06	5.51	53.16	2.92	2.65	4.85	4.59
CV		4.92	16.35	1.94	16.35	4.52	17.83	1.96	2.27	4.90	3.28

Data are mean values of n = 3 and are significantly different by the LSD test ( $p < 0.05$ ). <sup>\*\*</sup> and <sup>\*</sup> are significant at the 0.01 and 0.05 levels. Total flavonoid content in the leaf and root samples (TFL, TFR), total phenol content in the leaf and root samples (TPL, TPR), DPPH in the leaf and root samples (DPPHL, DPPHR), total antioxidant capacity in the leaf and root samples (TACL, TACR), and PFARAP assay in the leaf and root samples (PFRAPL, PFRAPR). NAA: 1-Naphthaleneacetic acid, BAP: 6-Benzylaminopurine, 2,4-D: 2,4-Dichlorophenoxyacetic acid.



**Fig. 3.** Total flavonoid content (A) and total phenol content (B) of *C. roseus* L. var. 'Ocellatus' on MS medium supplemented with different PGRs. Data are mean values of  $n = 3$ . Mean values followed by different letters indicate they are significantly different by the LSD test ( $p < 0.05$ ). NAA: 1-Naphthaleneacetic acid, BAP: 6-Benzylaminopurine, 2,4-D: 2,4-Dichlorophenoxyacetic acid.

### Antioxidant capacity

All three antioxidant assays revealed significant differences among the treatments in both leaf and root samples (Table 3).

Regarding DPPH, the highest radical scavenging was 74.21% ( $1 \text{ mg L}^{-1}$  NAA +  $0.5 \text{ mg L}^{-1}$  2,4-D) and 78.08% ( $1 \text{ mg L}^{-1}$  NAA +  $0.5 \text{ mg L}^{-1}$  2,4-D) in the leaf and root samples, respectively (Fig. 4A). The results varied from 74.21 to 65.66 in the leaves and from 78.08 to 38.34 in the roots.


The TAC assay indicated that maximum scavenging (89.33% and 87.9%) occurred in response to  $1 \text{ mg L}^{-1}$  NAA +  $1 \text{ mg L}^{-1}$  BAP and  $1 \text{ mg L}^{-1}$  NAA +  $1 \text{ mg L}^{-1}$  BAP +  $0.5 \text{ mg L}^{-1}$  2,4-D in the leaf and root samples, respectively (Fig. 4B). The comparison of mean values showed that the reducing power activity in

the leaves and root samples were affected by the  $0.5 \text{ mg L}^{-1}$  2,4-D +  $1 \text{ mg L}^{-1}$  BAP and  $1 \text{ mg L}^{-1}$  NAA +  $1 \text{ mg L}^{-1}$  BAP +  $0.5 \text{ mg L}^{-1}$  2,4-D treatments, thus having the highest significance level (62.34 and 62.68 in the leaves and roots, respectively) (Fig. 4C).

### Pearson's correlation coefficient

Pearson's correlation coefficient of TAC, TPC, DPPH, TAC, and RP from the roots were analyzed and the results indicated that a positive significant correlation was observed (Table 4). The highest correlation between phytochemical traits and antioxidant activity assays was related to the TPC with TAC ( $R^2 = 0.849$ ) and DPPH ( $R^2 = 0.845$ ). The correlation coefficient between DPPH and TAC was 0.966.





**Fig. 4.** Antioxidant assays: DPPH assay (A), TAC assay (B), and reducing power assay (C) for *C. roseus* L. var. 'Ocellatus' on MS medium supplemented with different PGRs. Data are mean values of  $n = 3$ . Mean values followed by different letters indicate they are significantly different by the LSD test ( $p < 0.05$ ). NAA: 1-Naphthaleneacetic acid, BAP: 6-Benzylaminopurine, 2,4-D: 2,4-Dichlorophenoxyacetic acid.

## Discussion

Fazal et al. (2020) reported that in the culture medium of *Prunella vulgaris* augmented with various NAA concentrations, maximum biomass production was detected, and NAA was the best treatment for fresh biomass production. As a root

regulator substance, auxin is suitable for controlling root development and its anatomy (Olatunji et al., 2017; Overvoorde et al., 2010), thus offering to encourage lateral root development, biosynthesis, polar transport, and signal transduction of auxin are crucial (Sun et al., 2017). Cytokinin is another central regulator of lateral root development

(Laplaze et al., 2007; Marhavý et al., 2011; Marhavý et al., 2014; Moreira et al., 2013). Several reports indicated the increasing role of combined auxins and cytokinins in fresh and dry root masses from *in vitro* plants (Danova et al., 2018). Morphogenesis of *Artemisia alba in vitro* plants was affected by exogenous cytokinins. According to Purohit et al. (2020) and Bakhshipour et al. (2019), auxin and cytokinins are suitable for plant growth

enhancement. Cytokinin via scavenging pin-formed (PIN) proteins leads to intervening with polar auxin transport and regulates lateral root development (Marhavý et al., 2011; Marhavý et al., 2014). Additional PGRs directly or indirectly contribute to cell fate decisions and cooperate differentially in lateral root development (Chandler and Werr, 2015; Schaller et al., 2015).

**Table 4.** Pearson's correlation coefficient of TFC, TPC, DPPH, TAC, and RP from the roots.

	Total flavonoid content	Total phenol content	DPPH assay	TAC assay	Reducing power assay
Total flavonoid content	1	0.389*	0.556**	0.633***	0.434*
Total phenol content		1	0.845***	0.849***	0.505*
DPPH assay			1	0.966***	0.551**
TAC assay				1	0.615**
Reducing power assay					1

Total flavonoid content (TFC), total phenol content (TPC), total antioxidant capacity (TAC), and DPPH (1,1-diphenyl-2-picrylhydrazyl) for the radical scavenging assay. \*, \*\* and ns indicate significance at the 0.05 and 0.01 levels and non-significance, respectively.

Kulus (2020) revealed that IAA (at 1 mg L<sup>-1</sup>) stimulated the synthesis of chlorophyll a and carotenoids in *Lamprocapnos spectabilis*, and similar results were obtained in previous research (Sarami et al., 2017), indicating that IAA was more effective than BAP on photosynthetic pigments. Changes in the chlorophyll content by PGRs could be related to growth rate and primary and secondary metabolic activities (Lichtenthaler, 1987). Kaviani (2014) and Parsaeimehr et al. (2010) reported the positive effects of auxin-cytokinin, separately or in combinations, on increasing the photosynthetic pigment content of *Ephedra*, *Eustoma grandiflorum* L. and *Saccharum officinarum* L. Furthermore, auxin-cytokinin revealed an increased cell division rate and chlorophyll content by the decrease in different enzymatic functions involved in chlorophyll biosynthesis. The involvement of various processes in the metabolism of chlorophyll and the multidimensionality of PGRs impacts is the reason for variable results (Bakhshipour et al., 2019).

These results were confirmed by a previous report (Fazal et al., 2020). Flavonoids are one of the largest groups of secondary metabolites belonging to polyphenols and play different roles in treatments by having a strong ability to degrade reactive oxygen species (Matkowski, 2008). In soybean (*Glycine max* L. Merr) plants, associations were studied between PGR treatments and the expression level of different genes involved in the biosynthesis of total phenolic, flavonoid, and isoflavones. Results showed that the 6-BA treatment caused the highest expression, followed by NAA and GA3 treatments (Guo et al., 2018). PGRs noticeably influence the production, distribution, and

accumulation of secondary metabolites, such as phenolics, and enhance the expression levels of associated genes (Radic, 2016). PGRs, especially a combination of applied auxin and cytokinins, can lead to an accumulation of flavonoids in cell culture. Their roles in the biosynthesis of flavonoids are well documented (Bota and Deliu, 2015; El-Shafey et al., 2016; Ji et al., 2015; Murthy et al., 2014). PGRs have shown significant effects on antioxidant scavenging, TPC, and TFC of micro-propagated *Vaccinium arctostaphylos* L. plants, and cytokinins treatments improved the antioxidant activity of *in vitro*-grown plants (Bakhshipour et al., 2019). Moreover, the extract of *in vitro*-grown *Coleonema pulchellum* L. plants showed high scavenging activity against DPPH radical (Baskaran et al., 2014). Another study that emphasized the roles of PGRs in the antioxidant activity of *in vitro*-grown plants was reported by Abbasi et al. (2019).

In plants, synergistic or additive functions between several growth hormones, such as auxins and cytokinins, by producing ethylene followed by an induced systemic resistance (ISR) mechanism have led to the appearance of total antioxidant capacity (Cary et al., 1995; Heinrich 2008; Van Wees et al., 2000; Woeste et al., 1999). Thus, growth hormones in the medium might affect the cells to generate ISR (Mustafa and Verpoorte, 2007) and increase antioxidant activity. Biochemical pathways and physiological processes in plants can be influenced by PGRs as they function. This signaling component can change plant metabolism and bioactive compounds (Mitra et al., 2016).

Aryal et al. (2019), Asem et al. (2020), Gan et al. (2017),

and Kainama et al. (2020) reported similar results about the presence of a strong correlation between TPC and different antioxidant activity assays. They suggested that bioactive compounds such as TPC and TFC play a scavenging role and exhibit antioxidant activity.

## Conclusion

An increase in demand for medicinal plants in recent years has led to the use of biotechnology techniques for the mass production of these plants in a uniform manner. Tissue culture techniques with ideal conditions for the micro-propagation of medicinal plants have excelled in producing these plants. An important and effective factor for *in vitro* production of medicinal plants is the use of PGRs or phytohormones. In addition to creating suitable growing conditions for plant production, these substances improve the phytochemical properties of medicinal plants. Here, the positive role of PGRs alone or in combination with each other enhanced the morphological, physiological, phytochemical properties, and antioxidant activity in the medicinal plant *Catharanthus roseus* L. Collectively, 1 mg L<sup>-1</sup> NAA alone or in combination with 1 mg L<sup>-1</sup> BAP improved the morphological attributes. Regarding phytochemical properties, 0.5 mg L<sup>-1</sup> 2,4-D treatments alone or combined with 1 mg L<sup>-1</sup> BAP resulted in optimal outcomes.

## Conflict of Interest

The authors indicate no conflict of interest in this work.

## References

- Abbasi BH, Siddiquah A, Tungmunthum D, Bose S, Younas M, Garros L, Drouet S, Giglioli-Guivarc'h N, Hano C. 2019. *Isodon rugosus* (Wall. ex Benth.) codd *in vitro* cultures: Establishment, phytochemical characterization and *in vitro* antioxidant and anti-aging activities. International Journal of Molecular Sciences 20, 452.
- Anand S. 2010. Various approaches for secondary metabolite production through plant tissue culture. Pharmacia 1, 1-7.
- Arnon DI. 1949. Copper enzymes in isolated chloroplasts polyphenol oxidase in *Beta vulgaris*. Plant Physiology 24, 1-15.
- Aryal S, Baniya MK, Danekhu K, Kunwar P, Gurung R, Koirala N. 2019. Total phenolic content, flavonoid content and antioxidant potential of wild vegetables from Western Nepal. Plants 8, 96.
- Asem N, Abdul Gapar NA, Abd Hapit NH, Omar EA. 2020. Correlation between total phenolic and flavonoid contents with antioxidant activity of Malaysian stingless bee propolis extract. Journal of Apicultural Research 59, 437-442.
- Bakhshpour M, Mafakheri M, Kordrostami M, Zakir A, Rahimi N, Feizi F, Mohseni M. 2019. *In vitro* multiplication, genetic fidelity and phytochemical potentials of *Vaccinium arctostaphylos* L.: an endangered medicinal plant. Industrial Crops and Products 141, 111812.
- Bakrudeen A, Subha Shanthy G, Gouthaman T, Kavitha M, Rao M. 2011. *In vitro* micropropagation of *Catharanthus roseus* – an anticancer medicinal plant. Acta Botanica Hungarica 53(1-2), 20.
- Baskaran P, Moyo M, Van Staden J. 2014. In vitro plant regeneration, phenolic compound production and pharmacological activities of *Coleonema pulchellum*. South African Journal of Botany 90, 74-79.
- Bota C, Deliu C. 2015. Effect of plant growth regulators on the production of flavonoids by cell suspension cultures of *Digitalis lanata*. Farmacia 63, 716-719.
- Cary AJ, Liu W, Howell SH. 1995. Cytokinin action is coupled to ethylene in its effects on the inhibition of root and hypocotyl elongation in *Arabidopsis thaliana* seedlings. Plant Physiology 107, 1075-1082.
- Chandler JW, and Werr W. 2015. Cytokinin–auxin crosstalk in cell type specification. Trends in Plant Science 20, 291-300.
- Chen TW, Yeh DC, Chao TY, Lin CH, Chow LW, Chang DY, Hsieh YY, Huang SM, Cheng AL, Lu YS, Consortium TBC. 2018. A phase I/II study of the combination of lapatinib and oral vinorelbine in HER2-positive metastatic breast cancer. Japanese Journal of Clinical Oncology 48 (3), 242–247.
- Classic Murashige T, and Skoog F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiologia Plantarum 15, 473-497.
- Danova K, Motyka V, Todorova M, Trendafilova A, Krumova S, Dobrev P, Andreeva T, Oreshkova T, Taneva S, Evstatieva L. 2018. Effect of cytokinin and auxin treatments on morphogenesis, terpenoid biosynthesis, photosystem structural organization, and endogenous isoprenoid cytokinin profile in *Artemisia alba* Turra *in vitro*. Journal of Plant Growth Regulation 37, 403-418.
- Derbassi NB, Pedrosa MC, Heleno S, Carocho M, Ferreira IC, Barros L. 2022. Plant volatiles: using scented molecules as food additives. Trends in Food Science and Technology.
- El-Shafey NM, Ahmed ES, Sayed M, Hammouda O, Khodary S. 2016. Effect of growth regulators, carbohydrates and antioxidant compounds on biomass, flavonoid accumulation and enzyme activity in callus cultures of *Rumex vesicarius* L. Egyptian Journal of Botany 56, 595-612.
- Fazal H, Abbasi BH, Ahmad N, Noreen B, Shah J, Ma D, Chuanliang L, Akbar F, Uddin MN, Khan H. 2020. Biosynthesis of antioxidative enzymes and polyphenolics content in calli cultures of *Prunella vulgaris* L. in response to auxins and cytokinins. Artificial Cells, Nanomedicine, and Biotechnology 48, 893-902.
- Gajalakshmi S, Vijayalakshmi S, Devi RV. 2013. Pharmacological activities of *Catharanthus roseus*: a perspective review. International Journal of Pharma and Bio Sciences 4, 431-439.
- Gan J, Feng Y, He Z, Li X, Zhang H. 2017. Correlations between antioxidant activity and alkaloids and phenols of Maca (*Lepidium meyenii*). Journal of Food Quality 2017.
- Guo X, Wang L, Chang X, Li Q, Abbasi AM. 2018. Influence of plant growth regulators on key-coding genes expression associated with phytochemicals biosynthesis and antioxidant activity in soybean (*Glycine max* (L.) Merr) sprouts. International Journal of Food Science & Technology, 1–9.
- Hatano T, Kagawa H, Yasuhara T, Okuda T. 1988. Two new flavonoids and other constituents in licorice root: their

- relative astringency and radical scavenging effects. *Chemical and Pharmaceutical Bulletin* 36, 2090-2097.
- Heinrich M. 2008. Ethnopharmacy and natural product research—multidisciplinary opportunities for research in the metabolomic age. *Phytochemistry Letters* 1, 1-5.
- Hernández-Domínguez E, Campos-Tamayo F, Vázquez-Flota F. 2004. Vindoline synthesis in *in vitro* shoot cultures of *Catharanthus roseus*. *Biotechnology Letters* 26(8), 671–674.
- Ji XH, Wang YT, Zhang R, Wu SJ, An MM, Li M, Wang CZ, Chen XL, Zhang YM, Chen XS. 2015. Effect of auxin, cytokinin and nitrogen on anthocyanin biosynthesis in callus cultures of red-fleshed apple (*Malus sieversii* f. *niedzwetzkyana*). *Plant Cell, Tissue and Organ Culture (PCTOC)* 120, 325-337.
- Kabesh K, Senthilkumar P, Ragunathan R, Kumar RR. 2015. Phytochemical analysis of *Catharanthus roseus* plant extract and its antimicrobial activity. *International Journal of Pure & Applied Bioscience* 3, 162-172.
- Kähkönen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS, Heinonen M. 1999. Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agricultural and Food Chemistry* 47, 3954-3962.
- Kainama H, Fatmawati S, Santoso M, Papilaya PM, Ersam T. 2020. The relationship of free radical scavenging and total phenolic and flavonoid contents of *Garcinia lasoar* PAM. *Pharmaceutical Chemistry Journal* 53, 1151-1157.
- Kaviani B. 2014. Micropropagation of ten weeks (*Matthiola incana*) and lisianthus (*Eustoma grandiflorum*) (two ornamental plants) by using kinetin (Kin), naphthalene acetic acid (NAA) and 2, 4–dichlorophenoxyacetic acid (2, 4–D). *Acta Scientiarum Polonorum Hortorum Cultus* 13, 141-154.
- Kulus D. 2020. Influence of growth regulators on the development, quality, and physiological state of *in vitro*-propagated *Lamprocapnos spectabilis* (L.) Fukuhara. *In Vitro Cellular & Developmental Biology-Plant* 56, 447-457.
- Laplaze L, Benkova E, Casimiro I, Maes L, Vanneste S, Swarup R, Weijers D, Calvo V, Parizot B, Herrera-Rodriguez MB. 2007. Cytokinins act directly on lateral root founder cells to inhibit root initiation. *Plant Cell* 19, 3889-3900.
- Lichtenthaler H.K. 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes, *Methods in Enzymology*. Elsevier, pp. 350-382.
- Marcone C, Ragozzino A, Seemuller E. 1997. Dodder transmission of alder yellows phytoplasma to the experimental host *Catharanthus roseus* (periwinkle). *European Journal of Forest Pathology* 27, 347-350.
- Marhavý P, Bielach A, Abas L, Abuzeineh A, Duclercq J, Tanaka H, Pařezová M, Petrášek J, Friml J, Kleine-Vehn J. 2011. Cytokinin modulates endocytic trafficking of PIN1 auxin efflux carrier to control plant organogenesis. *Developmental Cell* 21, 796-804.
- Marhavý P, Duclercq J, Weller B, Feraru E, Bielach A, Offringa R, Friml J, Schwechheimer C, Murphy A, Benková E. 2014. Cytokinin controls polarity of PIN1-dependent auxin transport during lateral root organogenesis. *Current Biology* 24, 1031-1037.
- Matkowski A. 2008. Plant *in vitro* culture for the production of antioxidants—a review. *Biotechnology Advances* 26, 548-560.
- Mitra D, Sharma K, Uniyal N, Chauhan A, Sarkar P. 2016. Study on plant hormone (indole-3-acetic acid) producing level and other plant growth promotion ability (pgpa) by *Asparagus racemosus* rhizobacteria. *Journal of Chemical and Pharmaceutical Research* 8, 995-1002.
- Montagna E, Palazzo A, Maisonneuve P, Canello G, Iorfida M, Sciandivasci A, Esposito A, Cardillo A, Mazza M, Munzone E, Lai A, Goldhirsch A, Colleoni M. 2017. Safety and efficacy study of metronomic vinorelbine, cyclophosphamide plus capecitabine in metastatic breast cancer: a phase II trial. *Cancer Letters* 400, 276–281.
- Moreira S, Bishopp A, Carvalho H, Campilho A. 2013. AHP6 inhibits cytokinin signaling to regulate the orientation of pericycle cell division during lateral root initiation. *PLoS One* 8, e56370.
- Murthy HN, Lee EJ, Paek KY. 2014. Production of secondary metabolites from cell and organ cultures: strategies and approaches for biomass improvement and metabolite accumulation. *Plant Cell, Tissue and Organ Culture (PCTOC)* 118, 1-16.
- Mustafa NR, Verpoorte R. 2007. Phenolic compounds in *Catharanthus roseus*. *Phytochemistry Reviews* 6, 243-258.
- Ochatt S, Alan A, Bhattacharya A, Hano C, Kiselev K, Marconi P, Otoni W, Park S, Tang K, Weathers P. 2022. Secondary metabolites: a boon from plants, the best chemist in nature: preface from the editors. *Plant Cell, Tissue and Organ Culture (PCTOC)* 1-6.
- Olatunji D, Geelen D, Verstraeten I. 2017. Control of endogenous auxin levels in plant root development. *International Journal of Molecular Sciences* 18, 2587.
- Overvoorde P, Fukaki H, Beeckman T. 2010. Auxin control of root development. *Cold Spring Harbor Perspectives in Biology* 2, a001537.
- Oyaizu M. 1986. Studies on products of browning reaction antioxidative activities of products of browning reaction prepared from glucosamine. *The Japanese Journal of Nutrition and Dietetics* 44, 307-315.
- Pandey-Rai S, Mallavarapu GR, Naqvi A, Yadav A, Rai SK, Srivastava S, Singh D, Mishra R, Kumar S. 2006. Volatile components of leaves and flowers of periwinkle *Catharanthus roseus* (L.) G. Don from New Delhi. *Flavour and Fragrance Journal* 21, 427-430.
- Parsaeimehr A, Sargsyan E, Javidnia K. 2010. A comparative study of the antibacterial, antifungal and antioxidant activity and total content of phenolic compounds of cell cultures and wild plants of three endemic species of Ephedra. *Molecules* 15, 1668-1678.
- Pereira DM, Ferreres F, Oliveira J, Valentão P, Andrade PB, Sottomayor M. 2009. Targeted metabolite analysis of *Catharanthus roseus* and its biological potential. *Food and Chemical Toxicology* 47, 1349-1354.
- Prieto P, Pineda M, Aguilar M. 1999. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. *Analytical Biochemistry* 269, 337-341.
- Purohit S, Joshi K, Rawat V, Bhatt ID, Nandi SK. 2020. Efficient plant regeneration through callus in *Zanthoxylum armatum* DC: an endangered medicinal plant of the Indian Himalayan region. *Plant Biosystems - an International Journal Dealing*

with all Aspects of Plant Biology 154, 288-294.

Radic S. 2016. Influence of pH and plant growth regulators on secondary metabolite production and antioxidant activity of *Stevia rebaudiana* (Bert). *Periodicum Biologorum* 118, 9–19.

Roberts J.A. 2012. Plant growth regulators. Springer Science & Business Media.

Sandhya M, Deepti L, Bhakti D, Ravindra M, Pranay S, Gauri A, Bansod I, Asmit H. 2016. Effect of growth regulator combination on *in-vitro* regeneration of *Catharanthus roseus*. *International Journal of Life Science* 6, 1-4.

Sarami R, Omidi H, Bostani A. 2017. The effect of auxin and cytokinin on some morpho-physiological and germination characteristics of *Stevia (Stevia rebaudiana Bertonii)* seeds. *Iranian Journal of Seed Research* 3.

Schaller GE, Bishopp A, Kieber JJ. 2015. The yin-yang of hormones: cytokinin and auxin interactions in plant development. *Plant Cell* 27, 44-63.

Singleton VL, Rossi JA. 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture* 16, 144-158.

Smetanska I. 2008. Production of secondary metabolites using plant cell cultures. *Food Biotechnology*, 187-228.

Sun CH, Yu JQ, Hu DG. 2017. Nitrate: a crucial signal during lateral roots development. *Frontiers in Plant Science* 8, 485.

Van Wees SC, De Swart EA, Van Pelt JA, Van Loon LC, Pieterse CM. 2000. Enhancement of induced disease resistance by simultaneous activation of salicylate- and jasmonate-dependent defense pathways in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences* 97, 8711-8716.

Woeste KE, Vogel JP, Kieber JJ. 1999. Factors regulating ethylene biosynthesis in etiolated *Arabidopsis thaliana* seedlings. *Physiologia Plantarum* 105, 478-484.

Yamamura J, Masuda N, Yamamoto D, Tsuyuki S, Yamaguchi M, Tanaka S, Tsurutani J, Tokunaga S, Yoshidome K, Mizutani M, Aono T, Ooe A, Tanino H, Matsunami N, Yasojima H, Nakayama T, Nishida Y. 2017. Gemcitabine and vinorelbine combination chemotherapy in taxane-pretreated patients with metastatic breast cancer: a phase II study of the Kinki Multidisciplinary Breast Oncology Group (KMBOG) 1015. *Chemotherapy* 62 (5), 307–313.

Yong JW, Ge L, Ng YF, Tan SN. 2009. The chemical composition and biological properties of coconut (*Cocos nucifera* L.) water. *Molecules* 14, 5144-5164.