



Influence of Tryptophan and Glycine Acids on Growth and Productivity of Fennel under Different Organic Fertilizer Levels

Ahmad Abdalazez Zatimeh*, Ahmad Hussin Al-Fraihat*

Department of Nutrition and Food Processing, Al-Huson University College, Al-Balqa Applied University Al-Salt 19117, Jordan

ARTICLE INFO

Article history:

Received: 24 May 2024,
Received in revised form: 14 August 2024,
Accepted: 24 August 2024

Article type:

Research paper

Keywords:

Farmyard manure,
Glycine,
Growth,
Tryptophan,
Volatile oil
Yield

COPYRIGHT

© 2025 The author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other medium is permitted, provided the original author(s) and source are cited, in accordance with accepted academic practice. No permission is required from the authors or the publishers.

ABSTRACT

This research aimed to improve the growth, productivity, and volatile oil production of fennel (*Foeniculum vulgare*, Mill.) plants. An experiment was conducted to evaluate the effects of farmyard manure (FYM) and amino acid mixtures (tryptophan and glycine) at different concentrations on fennel plants and their interaction parameters. The highest plant height was recorded with the application of 40 m³ ha⁻¹ FYM. Significant effects due to the applied organics were observed on the number of branches plant⁻¹, total dry weight plant⁻¹, number of umbels plant⁻¹, fruit yield, and volatile oil yield ha⁻¹. When compared with the abundance of other amino acids under study, fennel plants sprayed with 300 ppm T + glycine showed an increase in the above parameters. The best productivity and volatile oil was achieved in fennel plants grown with organic fertilizer (40 m³ ha⁻¹), including foliar spray of amino acids at a concentration of 300 ppm. The gas chromatography (GC/MS) analysis of fennel oil showed that the main compounds were estragole, D-limonene, L-fenchone, and anethole, which had the highest proportions of volatile oil compositions compared to other compounds. This study concluded that using organic fertilizers with foliar application and amino acids had positive effects on fennel growth, productivity, and volatile oil compounds.

Introduction

Foeniculum vulgare Mill., commonly referred to as fennel, is a medicinal and aromatic plant belonging to the family Umbelliferae (Apiaceae). Various parts of the fennel plant, including its leaves, stem, root, and particularly the fruits, possess aromatic properties and have been utilized in numerous applications (Badgujar et al.,

2014). The chemical constituents of fennel contribute to its medicinal properties (Kooti et al., 2015). Globally, fennel is employed to treat a wide array of ailments, including fever, flatulence, insomnia, renal disorders, cancer, stomachaches, liver discomfort, and abdominal pain (Choi and Hwang, 2004; Rather et al., 2012).

Plant nutrients play a crucial role in enhancing

*Corresponding author's email: ahmad@bau.edu.jo , azatimeh@bau.edu.jo

plant productivity. Organic manure, when added to soil, is broken down by microbes into minerals, releasing nutrients that improve the soil's cation exchange capacity and water retention (Kramer and Boyer, 1995). The addition of farmyard manure (FYM) enhances the chemical and physical properties of soils. Due to its organic matter content, comprising macro- and micro-elements, amino acids, sugars, and organic acids, FYM promotes soil fertility (Hamza et al., 2006). Furthermore, Ibrahim et al. (2006) demonstrated that applying FYM at a rate of 16 m³ per feddan increased vegetative growth and fruit yield per feddan in both fennel and caraway plants. Similarly, Abdelbaky et al. (2023) reported that among the seventeen identified components of fennel fruit essential oil, fenchone, trans-anethole, limonene, and estragole were highly abundant in FYM-treated plants. Al-Fraihat et al. (2011) indicated that the application of FYM resulted in maximum productivity in marjoram plants. More recently, Al-Fraihat et al. (2023a) found that high rates of organic fertilizers improved the growth, yield, and oil production of black cumin (*Nigella sativa* L.) and coriander (*Coriandrum sativum* L.).

Amino acids are vital in catalyzing secondary metabolic reactions in plants (Pratelli and Pilot, 2014). Tao et al. (2008) demonstrated that tryptophan is associated with the synthesis of camalexin, auxins, phenylpropanoids, phytoalexins, and other related natural compounds in plants. El-Awadi and Hassan (2010) observed significant increases in seed number per umbel, number of umbels per plant, and seed weight per umbel when the tryptophan concentration was set at 100 mg L⁻¹. Glycine also plays a key role in plant growth and chlorophyll synthesis, facilitating the absorption of Zn, Fe, Cu, and Mn (Ghasemi et al., 2013). With a molar mass of 75 g mol⁻¹, glycine is classified as a proteinogenic amino acid and a minor biological amino acid (Souri and Hatamian, 2019). The application of glycine at a concentration of 10 mg L⁻¹ in nutrient solutions has been shown to enhance plant growth and the nutritional quality of coriander (Al-Fraihat et al., 2023c; Mohammadipour and Souri, 2019). Similarly, Ali and Sallam (2022) demonstrated that using glycine or tryptophan at a high concentration (150 ppm) significantly improved the essential oil percentage in caraway compared to the control. Al-Fraihat et al. (2023b) further indicated that the application of amino acids (L-tryptophan and glutamine) resulted in enhanced growth of rosemary (*Rosmarinus officinalis* L.).

This study aimed to evaluate the effects of varying farmyard manure rates and the concentration of

amino acids (tryptophan and glycine) on the growth, fruit yield components, and essential oil production of fennel plants (*Foeniculum vulgare* Mill.) (Shahhoseini et al., 2020).

Material and Methods

A field experiment was conducted over two consecutive seasons, 2020–2021 and 2021–2022, at the experimental farm in the Ajloun region of Jordan. The aim was to investigate the effects of organic fertilizers, specifically farmyard manure (applied at rates of 0, 20, 30, and 40 m³ ha⁻¹), on fennel growth, yield, and volatile oil production. Additionally, the experiment assessed the impact of amino acid mixtures, such as tryptophan and glycine (applied at concentrations of 0, 100, 200, and 300 parts per million), as well as their interactions on these variables.

Before planting, the physical and chemical properties of the experimental soil were evaluated using methods outlined by Black (1965) and Page et al. (1982). Soil samples were taken from three random locations at a depth of 0–30 cm. The soil analysis was conducted at the National Center for Agricultural Research, Department of Laboratories in Jordan. The results of the soil analysis are presented in Table 1, followed by the nutritional composition of the analyzed fennel fruits, as shown in Table 2.

Experimental design and tested treatments

This experiment followed a randomized complete block design (RCBD) with three replicates, using a split-plot distribution. The main plots were farmyard manure rates (0, 20, 30 and 40 m³ ha⁻¹). A mixture of amino acids tryptophan and glycine at concentrations (0, 100, 200, and 300 ppm) were assigned as sub-plot. Fennel fruits were cultivated on 5 November of the bathing season. Amino acid treatments were sprayed at regular interval times of 45, 60, and 90 d from the sowing date.

Each sub-plot 3.0 × 3.0 m contained 5 rows each one 60 cm apart. The spacing between plants was 30 cm. After 33 d following the initial planting, the plants underwent thinning, producing two plants hill⁻¹. Data recorded in the first week of May included plant height (cm), number of branches plant⁻¹, fresh and dry weights (g). It also included number of umbels plant⁻¹, fruit yield plant⁻¹ (g), and fruit yield ha⁻¹ (kg). All of them in addition to volatile oil percentage were calculated in line with to Guenther (1961). Then, volatile oil yield plant⁻¹ (mL), and volatile oil yield ha⁻¹ (L) were computed. Before planting, mature cattle manure was put into the soil and mixed in. This was done to get the soil prepared for farming.

Table 1. Experimental soil analysis of the research area.

pH	EC (ds m ⁻¹)	ppm		Mg L ⁻¹				%			Texture	
		K	Ca	Mg	Cl	Na	HCO ₃	O.M	Clay	Silt		Sand
8.38	0.52	473	2.82	3.18	3.4	0.8	1.6	3.4	52.0	29.3	18.7	Clay

Table 2. Chemical analysis of farmyard manure in 2020-2021 and 2021-2022.

Constituent	Farmyard manure	
	2020/2021	2021/2022
EC ds m ⁻¹ (1:2 ex.)	3.40	3.33
pH (1:2 w/v)	7.06	7.03
O.M%	38.3	37.2
O.C%	22.5	21.6
N%	3.22	3.36
C/N ratio	20.1	21.1
P%	0.95	0.96
K%	3.66	3.71
Fe ppm	6250	6119
Mn ppm	229	235
Zn ppm	269	273
Cu ppm	14.8	15.11

Volatile oil insulation

To extract volatile oil, fruits from each treatment were gathered over both seasons and weighed. Then, 100 g of each duplicate was handled for three h during hydrodistillation (HD) using a Clevenger-type apparatus (Clevenger, 1928). Volatile oil content was calculated as a relative percentage (v/w). Likewise, total EO was calculated as mL 100 plants⁻¹ using dry weight. VOs taken out from *A. abrotanum* were gathered during the two seasons of each handling and dried on anhydrous sodium sulfate for chemical identification.

Gas chromatography and mass spectrometry (GC-MS) analysis

The Agilent 8890 GC System was leveraged for the analysis of the samples. It was linked to an Agilent 5977B GC/MSD mass spectrometer, which was fitted with an HP-5MS fused silica capillary column (30 m, 0.25 mm i.d., 0.25 mm film thickness). The oven was initially intended to operate at 50 °C, with a 5 °C min⁻¹ increase in temperature from 50 to 220 °C. At a rate of 20 °C min⁻¹, the temperature was then raised once more, from 220 °C to 280 °C. The temperature was kept at 280 °C for five min. The carrier gas, helium, was made use of at a flow rate of 1.0 mL min⁻¹. A gas chromatograph (GC) was used to analyze 1 µL of the dissolved volatile oil (30 µL of essential oil mL⁻¹ of diethyl ether). The split ratio of the GC was set to 1:50. At injection, the temperature was

230 °C. Using electron impact (EI) mode, which has a scan range of m/z 39 to 500 amu, mass spectra were obtained at an energy of 70 electron volts (eV) (Araghi, 2019; Taghizadeh et al., 2021). An experimental method was used on a split-plot design with RCBD. Gomez and Gomez's (1984) methodology was followed in the evaluation of the data. The statistical software package, Statistics Version 9, was utilized to compare the derived means (Analytical Software, 2008).

Results

Plant growth parameters

The results presented in Table 3 demonstrate that fennel growth and development were significantly higher at all farmyard manure (FYM) application rates (20, 30, and 40 m³ ha⁻¹) in both seasons, compared to the control. The highest values for plant height, number of branches per plant, total fresh weight, and total dry weight per plant were observed at the highest FYM rate (40 m³ ha⁻¹) during the 2020–2021 and 2021–2022 seasons. The application of organic fertilizers not only provided essential nutrients for plant nutrition but also contributed to the long-term sustainability of agricultural ecosystems (Maçik et al., 2020; Shaji et al., 2021). Organic fertilizers, derived from natural sources, supply necessary nutrients to plants.

Similarly, Massoud et al. (2019) reported that the growth of *Petroselinum sativum* (as measured by plant height, fresh weight, and dry weight) was

enhanced with increasing rates of FYM at six tons per feddan, reaching maximum values. In contrast, Abd El-Kafie et al. (2020) found that coriander plants fertilized with organic fertilizers

exhibited greater plant height and branch number compared to those fertilized with synthetic fertilizers such as FYM.

Table 3. Impacts of the rate of farmyard manure, amino acids, and their combinations on growth parameters of *Foeniculum vulgare* in 2020-2021 and 2021-2022.

FYM	Growth parameters								
	Plant height (cm)		Number of branches plant ⁻¹		Total fresh weight plant ⁻¹ (g)		Total dry weight plant ⁻¹ (g)		
	1 st season	2 nd season	1 st season	2 nd season	1 st season	2 nd season	1 st season	2 nd season	
Effect of FYM rate (m³ ha⁻¹)									
Control	114.83	116.50	7.73	9.67	155.50	163.92	35.73	40.64	
20	118.58	121.67	9.13	11.31	163.17	170.92	38.15	43.27	
30	121.17	127.00	10.58	13.93	170.33	178.17	40.85	44.64	
40	128.00	132.17	12.90	16.13	177.00	186.58	43.27	46.96	
L.S.D. at 5%	1.45	2.09	0.32	0.65	1.51	1.56	0.35	0.74	
Effect of AM concentration (ppm)									
Control	117.58	121.83	9.23	11.98	163.75	172.17	38.52	42.88	
100	119.50	123.50	9.81	12.48	165.42	174.00	39.17	43.52	
200	121.92	125.58	10.38	12.92	167.25	175.58	39.18	44.13	
300	123.58	126.42	10.92	13.67	169.58	177.83	40.50	44.98	
L.S.D. at 5%	0.37	0.42	0.19	0.21	0.34	0.40	0.18	0.81	
Effect of combination between FYM and AM									
Control	Control	112.00	114.67	6.83	9.08	152.33	161.67	34.92	39.65
	100	113.67	115.67	7.50	9.42	154.00	163.00	35.42	40.25
	200	116.33	117.00	8.08	9.92	156.67	164.33	36.00	40.92
	300	117.33	118.67	8.50	10.25	159.00	166.67	36.58	41.73
20	Control	115.33	119.33	8.42	10.58	160.67	168.33	37.17	42.32
	100	118.00	121.33	8.92	11.08	162.33	170.00	37.67	42.92
	200	119.33	124.00	9.42	11.33	163.67	171.67	38.50	43.50
	300	121.67	122.00	9.75	12.25	166.00	173.67	39.25	44.33
30	Control	118.33	124.00	9.83	13.00	168.00	175.00	39.83	43.57
	100	119.67	126.33	10.50	13.58	169.67	177.00	40.42	44.50
	200	122.67	128.33	10.75	14.08	171.00	179.00	41.17	44.83
	300	124.00	129.33	11.25	15.08	172.67	181.67	42.00	45.67
40	Control	124.67	129.33	11.83	15.25	174.00	183.67	42.15	46.00
	100	126.67	130.67	12.33	15.83	175.67	186.00	43.17	46.42
	200	129.33	133.00	13.25	16.33	177.67	187.33	43.58	47.25
	300	131.33	135.67	14.17	17.08	180.67	189.33	44.17	48.17
L.S.D. at 5%	1.59	2.21	0.46	0.74	1.62	1.71	0.47	1.58	

FYM = Farmyard manure (20, 30 and 40 m³ ha⁻¹) and AM = Amino acids (100, 200, and 300 ppm).

According to Table 3, applying an amino acid mixture (tryptophan and glycine) at a high concentration (300 ppm) to fennel plants led to a significant improvement in plant height (123.58 and 126.42 cm). It also increased the number of branches plant⁻¹ (10.92 and 13.67 branches),

total fresh (169.58 and 177.83 g plant⁻¹), and dry weights (40.50 and 44.98 g plant⁻¹). They contrasted with the control and the other concentrations being investigated in the first and second seasons, respectively. Generally, increasing amino acid concentrations gradually

increased fennel growth parameters in both seasons. Moreover, certain amino acids applied topically may have positive impacts on vegetative development and yield (Sadak et al., 2015; Shams et al., 2016). Likewise, compared to the control, Soliman et al. (2023) indicated that spraying glycine at 200 and 400 ppm concentrations considerably boosted the vegetative growth of cumin plants, as measured by plant height, branch number plant⁻¹, and fresh and dry weights of cumin.

Table 3 showed that spraying fennel plants thrice season⁻¹ with amino acids at 300 ppm concentration + 40 m³ ha⁻¹ of FYM recorded increased values. These increased values in plant growth parameters contrasted to other composite coefficients used in study coefficients.

Plant height, branch count, and fresh and dry weights plant⁻¹ were all gradually increased in both seasons by increasing amino acid concentrations under any rate of fiber wasting management. As mentioned before, plant growth is generally enhanced by the use of amino acids and organic fertilizer separately. Yet, when combined, their impacts can be even more pronounced, producing the tallest fennel plants, the most branches plant⁻¹, and the heaviest fresh and dry weights plant⁻¹.

Yield and its constituents

Table 4 show that using FYM fertilization treatments at a high rate (40 m³ ha⁻¹) significantly gave the highest values in umbels number plant⁻¹ (29.42-33.25 umbels). It also the highest values in fruit yield fennel plant⁻¹ (39.33-42.50 g plant⁻¹) and fruit yield ha⁻¹ (3186.7-3400.0 kg ha⁻¹); being contrasted to control and the lowest rates in both seasons, respectively. Likewise, all FYM rates (20, 30, and 40 m³ ha⁻¹) significantly increased yield components of fennel (*Foeniculum vulgare* Mill) contrasted to control during the two seasons. Following El-Sayed et al. (2017), the use of 100% organic fertilizer resulted in the greatest umbel count, umbel diameter, and umbel number dill plant⁻¹. Moreover, when contrasted with other treatments for the dried ripe fruits, the 100% organic fertilizer treatment produced the finest results in terms of fruit weight (g plant⁻¹) and fruit output (t fed⁻¹). Kumar et al. (2019) showed the maximum values for several different order umbels, seed yield, and biological yield of *Daucus carota* plants. They were recorded with FYM at 20 t ha⁻¹ in both years.

Table 4 showed increases in several umbels plant⁻¹ (4.00-3.47 branches) and fruit yield plant⁻¹ (3.67-4.34 g plant⁻¹). They also show the recorded

increases in fruit yield ha⁻¹ (293.3-346.6 kg ha⁻¹). This happened when fennel plants fertilized with 40 m³ ha⁻¹ in the first and second season, respectively. All FYM fertilization rates significantly increased yield components of fennel in contrast to the control during the two seasons. Furthermore, increasing FYM rates gradually increased fennel yield parameters in both seasons.

Table 4 showed that all combination treatments between FYM fertilization at 40 m³ ha⁻¹ and a mixture of amino acids (tryptophan and glycine) caused significant augments. These augments exemplified the number of umbels fennel plant⁻¹, fruit yield plant⁻¹, and ha⁻¹, contrasted to unfertilized plants. Furthermore, the yield components of the fennel plant were increased because of all combination treatments between FYM fertilization rates and amino acids contrasted to that of FYM fertilization alone. Gendy et al. (2015) indicated that foliar application of 100 ppm Tryptophan resulted in a visible increase in yield components of fenugreek (number of pods plant⁻¹, seed yield plant⁻¹, and seed yield ha⁻¹). In addition, Lemma (2022) revealed that the highest fruit number plant⁻¹; dry sepals yield every ha, and biological yield every ha was recorded with 15 t of FYM ha⁻¹.

Volatile oil production

Table 5 clarified that FYM rate treatments significantly increased fennel volatile oil production in contrast to the control in 2020/2021 and 2021/2022. The highest values in *Foeniculum vulgare* volatile oil percentage (2.256-2.419%), volatile oil yield plant⁻¹ (0.900-1.029 mL plant⁻¹), and volatile oil yield ha⁻¹ (71.980-82.320 L ha⁻¹) were obtained from 40 m³ FYM ha⁻¹. These were in contrast to the control and the other rates during the study in the first and second seasons, respectively. Also, Kamal et al., 2023; El-Sayed et al., 2017 found that using 100% organic fertilizer resulted in an elevated proportion of dill essential oil. Also, it caused the highest essential oil content (mL plant⁻¹) and essential oil production (L ha⁻¹) when contrasted to other treatments. This was observed across two growing seasons using dried ripe fruits. In line with Kamal et al. (2023), organic fertilizer influenced coriander's volatile oil. Additionally, applying 20 m³ of compost feddan⁻¹ raised the proportion of volatile oil in both seasons, while unfertilized crops had the least significant values. Table 5 results show exposing fennel plants to amino acids mixture (tryptophan and glycine) concentrations significantly increased volatile oil percentage, volatile oil yield plant⁻¹ (mL), and ha

¹ (L), compared to the control (unsprayed plants) in 2020/2021 and 2021/2022 seasons. In addition, the highest values in fennel volatile oil production (2.149 and 2.318%, 0.794 and 0.925 mL plant⁻¹ as well as 63.483 and 73.996 L ha⁻¹) were produced in response to 300 ppm of amino

acids concentration contrasted to the control, with other concentrations during the study. Generally, a progressive increase in this concern was brought about by an increase in the concentrations of amino acids during both seasons.

Table 4. Influence of the concentration of amino acids, the rate of farmyard manure, and their combinations on the yield components of *Foeniculum vulgare* in the seasons of 2020-2021 and 2021-2022.

FYM	Yield components						
	Number of umbels plant ⁻¹		Fruit yield plant ⁻¹ (g)		Fruit yield ha ⁻¹ (kg)		
	1st season	2nd season	1st season	2nd Season	1st Season	2nd season	
Effect of FYM rate (m3 ha⁻¹)							
Control	23.75	27.08	29.50	32.33	2360.0	2586.7	
20	25.42	29.83	33.75	36.33	2700.0	2906.7	
30	27.42	31.33	36.67	39.67	2933.3	3173.3	
40	29.42	33.25	39.33	42.50	3186.7	3400.0	
L.S.D. at 5%	1.18	0.83	0.57	1.19	45.7	94.9	
Effect of AM concentration (ppm)							
Control	24.75	28.95	33.00	35.33	2640.0	2826.7	
100	25.67	29.58	34.50	37.33	2760.0	2986.7	
200	26.83	30.75	35.58	38.50	2846.7	3080.0	
300	28.75	32.42	36.67	39.67	2933.3	3173.3	
L.S.D. at 5%	0.06	0.36	0.23	0.44	18.6	35.1	
Effect of combination between FYM and AM							
Control	Control	22.00	25.33	28.00	30.33	2240.0	2426.7
	100	23.00	26.33	29.00	32.00	2320.0	2560.0
	200	24.00	27.33	30.00	33.00	2400.0	2640.0
	300	26.00	29.33	31.00	34.00	2480.0	2720.0
20	Control	23.67	28.00	32.00	34.00	2560.0	2720.0
	100	24.67	29.00	33.33	36.00	2666.7	2880.0
	200	25.67	30.33	34.33	37.00	2746.7	2960.0
	300	27.67	32.00	35.33	38.33	2826.7	3066.7
30	Control	25.67	30.00	34.67	37.33	2773.3	2986.7
	100	26.67	30.33	36.33	39.33	2906.7	3146.7
	200	27.67	31.67	37.33	40.33	2986.7	3226.7
	300	29.67	33.33	38.33	41.67	3066.7	3333.3
40	Control	27.67	31.67	37.33	39.67	2986.7	3173.3
	100	28.33	32.67	39.33	42.00	3146.7	3360.0
	200	30.00	33.67	40.67	43.67	3253.3	3493.3
	300	31.67	35.00	42.00	44.67	3360.0	3573.3
L.S.D. at 5%	1.21	1.03	0.70	1.40	55.7	112.3	

FYM = Farmyard manure (20, 30 and 40³ ha⁻¹) and AM = Amino acids (100, 200, and 300 ppm).

Table 5 revealed that increasing amino acid concentrations from 200 to 300 ppm under each FYM rate gradually increased fennel volatile oil percentage; yield plant⁻¹; yield feddan⁻¹ in both seasons. The highest values in this regard were

(2.337- 2.480%, 0.981-1.108 mL plant⁻¹ in conjunction with 78.490 and 88.630 L ha⁻¹). They were achieved by FYM at 40 m³ ha⁻¹ + mixture of amino acids at 300 ppm contrasted to other combinations under consideration in both

seasons, respectively.

In addition, fennel plant volatile oil yield increased as a percentage and plant⁻¹ and ha⁻¹ when FYM fertilization and different amino acid

concentrations were used alone. Together, these factors may have maximized the effects of the individual treatments, resulting in higher volatile oil yield plant⁻¹ and ha⁻¹.

Table 5. Influence of farmyard manure rate, amino acid concentration and their combinations on volatile oil production of *Foeniculum vulgare* during 2020/2021 and 2021/2022 seasons.

FYM	Yield components						
	Essential oil percentage		Essential oil yield plant ⁻¹ (mL)		Essential oil yield ha ⁻¹ (L)		
	1 st season	2 nd season	1 st Season	2 nd Season	1 st season	2 nd season	
Effect of FYM rate (m³ ha⁻¹)							
Control	1.885	2.048	0.557	0.663	44.538	53.044	
20	1.988	2.183	0.672	0.795	53.754	63.554	
30	2.121	2.322	0.779	0.922	62.312	73.730	
40	2.256	2.419	0.900	1.029	71.980	82.320	
L.S.D. at 5%	0.057	0.024	0.029	0.028	2.362	2.198	
Effect of AM concentration (ppm)							
Control	1.974	2.153	0.656	0.767	52.483	61.312	
100	2.036	2.224	0.708	0.836	56.652	66.846	
200	2.090	2.276	0.749	0.881	59.960	70.495	
300	2.149	2.318	0.794	0.925	63.483	73.996	
L.S.D. at 5%	0.053	0.011	0.019	0.013	1.541	1.029	
Effect of combination between FYM and AM							
Control	Control	1.813	1.940	0.508	0.589	40.623	47.080
	100	1.860	2.027	0.539	0.649	43.140	51.873
	200	1.907	2.090	0.572	0.690	45.770	55.183
	300	1.960	2.133	0.608	0.726	48.617	58.040
20	Control	1.890	2.083	0.605	0.709	48.393	56.680
	100	1.957	2.163	0.652	0.779	52.200	62.303
	200	2.020	2.223	0.694	0.822	55.530	65.787
	300	2.083	2.263	0.736	0.868	58.893	69.447
30	Control	2.023	2.227	0.702	0.831	56.143	66.477
	100	2.093	2.310	0.763	0.908	60.990	72.670
	200	2.150	2.353	0.802	0.949	64.183	75.907
	300	2.217	2.397	0.849	0.999	67.933	79.867
40	Control	2.170	2.363	0.810	0.938	64.773	75.010
	100	2.233	2.397	0.878	1.007	70.280	80.537
	200	2.283	2.437	0.929	1.064	74.357	85.103
	300	2.337	2.480	0.981	1.108	78.490	88.630
L.S.D. at 5%	0.108	0.031	0.044	0.035	3.552	2.819	

FYM = Farmyard manure (20, 30 and 40³ ha⁻¹) and AM = Amino acids (100, 200, and 300 ppm).

Volatile oil components

The results of fennel oil gas chromatographic/mass spectrometry (GC/MS) indicated that it contains twelve different components (Table 6). When comparing the

chemical compound values in the oil, we find that, in contrast to other compounds, the aromatic oil compounds with the largest percentages were enethral, limonene, phenol, and anethole.

Table 6. Influence of farmyard manure rate, amino acid concentration and their combinations on volatile oil components of *Foeniculum vulgare* (mixed samples of 2020/2021).

No	Component Name	R T	Treatments				
			control	FYM 30 m ³ ha ⁻¹ + AM 200 ppm	FYM 30 m ³ ha ⁻¹ + AM 300 ppm	FYM 40 m ³ ha ⁻¹ + AM 200 ppm	FYM 40 m ³ ha ⁻¹ + AM 300 ppm
1	<i>α</i> -Pinene	6.316	1.04	0.91	1.42	1.15	1.11
2	<i>β</i> -Phellandrene	7.281	0.24	0.30	0.31	0.29	0.33
3	<i>β</i> -Myrcene	7.691	0.30	0.35	0.41	0.36	0.43
4	<i>α</i> -Phellandrene	8.078	---	----	0.23	---	0.21
5	<i>p</i> -Cymene	8.609	0.30	0.28	0.37	0.33	0.39
6	D-Limonene	8.725	11.35	12.90	13.28	11.94	13.26
7	Eucalyptol	8.811	0.56	0.48	0.58	0.46	0.55
8	<i>trans</i> - <i>β</i> -Ocimene	8.933	0.61	0.68	0.83	0.75	0.86
9	<i>γ</i> -Terpinene	9.533	0.35	0.35	0.40	0.44	0.42
10	L-Fenchone	10.382	4.34	4.12	4.30	4.87	4.28
11	Estragole	13.559	78.3	76.27	75.50	77.54	75.44
12	Anethole	15.852	2.6	3.39	2.18	1.95	2.24
Number of vehicles selected			11	11	12	12	12
Total % of compounds identified			100	100	100	100	100

FYM = Farmyard manure (20, 30, and 40³ ha⁻¹) and AM = amino acids (100, 200, and 300 ppm).

The uppermost average was recorded in the Estragole compound (78.30) in control, followed by FYM at 40 m³ ha⁻¹ + 200 ppm amino acids (AM), which recorded (77.54). Then FYM at 30 at m³ ha⁻¹ + 200 ppm amino acids (AM), which was (76.27) followed. The highest average was for the compound D-Limonene (13.28) in FYM at 30 at m³ ha⁻¹ + 300 ppm amino acids. FYM at 30 at m³ ha⁻¹ + 200 ppm amino acids, which recorded (12.90), followed. The highest percentages were for the compound L-Fenchone (4.87) in treatment FYM at 40 m³ ha⁻¹ + 200 ppm amino acids. The highest relative to the anethole compound (3.39) in treatment FYM at 30 at m³ ha⁻¹ + 200 ppm amino acids (AM). This implies the treatments had a discernible impact on raising the concentrations of some of the fundamental fennel oil constituents. Additionally, the treatments clearly affect several chemicals. This complies with the findings of Mahfouz et al. (2007) that using fertilizers increased the proportion of oil and aromatic oil components. This is explained by the integration of the two forms of fertilization and how it affects the availability of every component. Nutrients for the plant and this outcome support the findings of (DARWISH et al., 2011) that fertilizers increase the amount of chemicals present in fennel plants.

Discussion

The registered results of this investigation demonstrated the potential of organic manures to increase dill chemical constituents, yield components, and rate of plant growth. Numerous researchers have noted the significant functions that organic manures play in biological and physiological processes, which may account for the increase in these earlier features. Franz (1973) showed that augmenting the content of organic matter in soils caused an increase in dehydrogenase activity. In addition, organic manure holds moisture and maintains sufficient pore spaces to permit good air circulation and excessive water drainage. Schachtschable (1979) and Bohn et al. (1985) concluded that organic matter is a main source of N, P, S and contains high content of B and Mo. Also, it is considered as a source of energy for Azotobacter growth. Organic manure minimizes the loss of nutrients by leaching (Saber, 1997) and it caused an increase in microbial activities in the root zone when supplied to the soil (Taiwo et al., 2002). Mashali (1997) demonstrated that the incorporation of organic manure in the soil improves the permeability of soil and releases carbon dioxide and certain organic acids during decomposition. Amino acids serve as fundamental components of

proteins and the precursors for several other compounds that play vital roles in plants. Amino acids participate in the production of other chemical molecules, including alkaloids, plant hormones enzymes, and terpenoids that regulate diverse plant functions (Glawischnig et al., 2000; Ibrahim et al., 2010). Soliman et al. (2023) concluded that using glycine at 200 and 400 ppm as a foliar spray on cumin plants significantly affected the volatile oil percentage, associated with volatile oil yield plant⁻¹ and ha⁻¹.

Tryptophan is a necessary amino acid that plants need for their growth because it plays a role in the production of natural auxins called indole acetic acid (IAA). It affects the yield of plants (Yassen et al., 2010). Also, many plants and horticultural crops, in particular, depend heavily on glycine for proper nutritional management. Because of its simple technique of synthetic production, there is interest in mixing it with minerals to make chelates, which will help plants absorb and transport nutrients more efficiently (Souri and Hatamian, 2019).

Conclusion

The findings suggest that 40 m³ of FYM acre⁻¹ in conjunction with a 300 ppm foliar spray of an amino acid combination is advised to enhance fennel plant growth, yield components, and volatile oil production.

Conflict of Interest

The authors indicate no conflict of interest in this work.

References

Abdelbaky AS, Mohamed AM, Abd El Mageed TA, Rady MM, Alshehri F, El Saadony MT, Abun Qamar SF, El Tarabily KA, Al Elwany OAA. 2023. Bio-organic fertilizers promote yield, chemical composition, and antioxidant and antimicrobial activities of essential oil in fennel (*Foeniculum vulgare*) seeds. *Scientific Reports* 13, 1–16.

AbdEL-Kafie O.M, Ghaly N.G, El-Banna H.Y, Mostafa A.A.H. 2020. Effect of fertilization and foliar application treatments on coriander plant (*Coriandrum sativum* L.). *Journal of Plant Production, Mansoura Univ.* 11 (12): 1375-1382.

Ali HMM, Sallam DA. 2022. Inducing the growth and flowering of the caraway (*Carum carvi*L.) plant. *Journal of Applied Biology and Biotechnology* 10 (4), 86–91.

Al-Fraihat AH, Al-Dalain SY, Al-Rawashdeh ZB, Abu-Darwish MS, Al-Tabbat JA. 2011. Effect of organic and biofertilizers on growth, herb yield, and volatile oil of marjoram plant grown in Ajloun region, Jordan. *Journal of Medicinal Plants Research* 5, 2822–2833.

Al-Fraihat AH, Al-Dalain SY, Zatimeh AA, Haddad MA.

2023a. Enhancing Rosemary (*Rosmarinus officinalis* L.) growth and volatile oil constituents grown under soil salinity stress by some amino acids. *Horticulturae* 9 (2), 252. <http://dx.doi.org/10.3390/horticulturae9020252>.

Al-Fraihat AH, Zatimeh AA, Al-Hrout HH, Al-Dalain SY, Mostipan M. 2023b. The interaction impact of compost and biostimulants on growth, yield and oil content of black cumin (*Nigella sativa* L.) plants. *Brazilian Journal of Biology = Revista Brasileira de Biologia* 83, e272957. PMID: 37610946.

Al-Fraihat AH, Zatimeh AA, Al-Dalain SY. 2023c. The efficiency of salicylic acid and poultry manure on growth and volatile oil production of *Coriandrum sativum* L. plants. *Brazilian Journal of Biology* 84, e276886. doi: 10.1590/1519-6984.276886. PMID: 37991061.

Analytical Software. 2008. *Statistics Version 9*. Analytical Software, Tallahassee, Florida, USA.

Araghi AM, Nemati H, Azizi M, Moshtaghi N, Shoor M, Hadian J. 2019. Assessment of phytochemical and agromorphological variability among different wild accessions of *Mentha longifolia* L. cultivated in field condition. *Industrial Crops and Products* 140, 111698.

Badgujar SB, Patel VV, Bandivdekar AH. 2014. *Foeniculum vulgare* Mill.: A review of its Botany, Phytochemistry, Pharmacology, Contemporary Application and Toxicology. *BioMed Research Journal* 1, 1–32.

Black CA. 1965. *Methods of Soil Analysis. Part 2. Agronomy Monograph No. 9*, ASA, Madison, Wisconsin, USA.

Bohn H.L., Meneal B.L., Connar G.A.O. 1985. *Soil Chemistry*. 2nd ed. Wiley Interscience Publication. John Wiley and Sons, New York.

Choi EM, Hwang JK. 2004. Anti-inflammatory, analgesic, and antioxidant activities of the fruit of *Foeniculum vulgare*. *Fitoterapia* 75, 557–565.

Clevenger JF. 1928. Apparatus for Volatile Oil Determination, Description of New Type. *The Journal of the American Pharmaceutical Association* 17, 345–349.

Darwish MA, Al-Fraihat A, Dalaeen S, Afifi F, Al-Tabbat J. 2011. Determination of essential oils and heavy metals accumulation in *Salvia officinalis* cultivated in three intra-row spacing in Ash-Shoubak, Jordan. *International Journal of Agriculture and Biology* 13, 981-985.

El-Awadi ME, Hassan EA. 2010. Physiological responses of fennel (*Foeniculum vulgare* Mill) plants to some growth substances. The effect of certain amino acids and a pyrimidine derivative. *Journal of American Science* 6 (7), 120–127.

El-Sayed AA, Darwish MA, Azoz SN, Abd-Alla ABM, Elsayed SIM. 2017. Effect of mineral, bio and organic fertilizers on productivity, essential oil composition and fruit anatomy of two dill cultivars (*Anethum graveolens* L.). *Middle East Journal of Applied Sciences*

7 (3), 532–550.

Franz G. 1973. Comparative investigations on the enzyme activity of some soil in Nordrhein Westfalen and Rhein landpflaz. *Pedobiologia* 13(6), 423. CC.F. Soil and Fertilizers. 37, 8.

Gendy ASH, Soliman AH, Mohammed NS. 2015. Response of growth, productivity and oil composition of fenugreek plants to foliar application of complete fertilizer, dry yeast and L-tryptophan under sandy soil conditions. *Current Science International* 4 (4), 736–749.

Ghasemi P, Sabet P, Aadal H, Hadi M, Jamshidi M, Golchin K, R. 2013. Application of Domino Theory to Justify and Prevent Accident Occurrence in Construction Sites. *IOSR Journal of Mechanical and Civil Engineering* 6 (2), 2320–2334. Retrieved from www.iosrjournals.org

Glawischnig E, Tomas A, Eisenreich W, Spiteller P, Bacher A, Gierl A. 2000. Auxin biosynthesis in maize kernels. *Plant Physiology* 12 (3), 1109–1120.

Gomez NK, Gomez AA. 1984. *Statistical Procedures for Agricultural Research*. 2nd Ed. John Wiley and Sons, New York, USA, 680.

Guenther E. 1961. *The Essential Oil*. D. Von Nostrand Company, New York. 1, 236.

Hamza AM, Massoud HY, Eid ME, Khater MR, El-Gamal SMA. 2006. Effect of farmyard manure (FYM) doses and different bio-fertilizers on vegetative growth, seed yield and active constituents of *Plantago ovata* Forsk plants. *Journal of Agricultural Science, Mansoura University* 32 (7), 5583–5600.

Ibrahim SM, El-Labban HM, Mohamed FI, Naga NM. 2006. Effect of organic manures and chemical fertilizers on *Foeniculum vulgare* Mill and *Carum carvi* L. *Bulletin of Pharmaceutical Sciences, Assiut University* 29 (1), 187–201.

Ibrahim SMM, Taha LS, Farahat MM. 2010. Influence of foliar application of pepton on growth, flowering and chemical composition of *Helichrysum bracteatum* plants under different irrigation intervals. *Ozean Journal of Applied Sciences* 3 (1), 143–155.

Kamal SA, Kandeel AM, Hassan SE, Radwan AS. 2023. Impact of planting dates and organic fertilization on growth, productivity and essential oil components of coriander (*Coriandrum sativum* L.) plant. *Egyptian Journal of Horticulture* 50 (1), 73–84.

Kramer PJ, Boyer JS. 1995. *Water Relations of Plants and Soils*. Academic Press, San Diego, USA.

Kooti W, Moradi M, Ali-Akbari S, Sharafi-Ahvazi N, Asadi-Samani M, Ashtary-Larky D. 2015. Therapeutic and pharmacological potential of *Foeniculum vulgare* Mill.: A review. *Journal of HerbMed Pharmacology* 4 (1), 1–9.

Kumar A, Batra VK, Panghal VS, Bhuker A, Kumar R. 2019. Influence of FYM and gypsum on growth and seed yield in carrot (*Daucus carota* L.) irrigated with high RSC water. *International Journal of Current*

Microbiology and Applied Sciences 8 (3), 337–345.

Lemma DT. 2022. Evaluation of yield and economic profitability of using NP and FYM for the production of roselle. *International Research Journal of Plant Science* 13 (4), 1–10.

Maçik M, Gryta A, Frac M. 2020. Bio-fertilizers in agriculture: An overview on concepts, strategies and effects on soil microorganisms. In: Sparks DL (ed) *Advances in Agronomy*. Academic Press, 31–87.

Mahfouz S, Sharaf Eldin MA. 2007. Effect of mineral vs. biofertilizer on growth, yield, and essential oil content of fennel (*Foeniculum vulgare* Mill). *International Agrophysics* 21 (4), 361–366.

Masada KHK, Ben Taarit M, Chahed T, Kchouk ME, Marzouk B. 2007. Changes on essential oil composition of coriander (*Coriandrum sativum* L.) fruits during three stages of maturity. *Food Chemistry* 102, 1131–1134.

Mashali A.M. 1997. FAO global network on integrated soil management for sustainable use of salt affected soils. *Proceedings of International Symposium on Sustainable Management of Salt Affected Soils in the Arid Ecosystem*, 21-26 September, Cairo, Egypt, 31-35.

Massoud HYA, Dawa KK, ElGamal SMA, Karkash SHA. 2019. Response of (*Petroselinum sativum* Hoffm.) to organic, bio-fertilizer and some foliar application. *Journal of Plant Production, Mansoura University* 10 (12), 1194–1161.

Mohammadipour N, Souri MK. 2019. Effects of different levels of glycine in the nutrient solution on the growth, nutrient composition, and antioxidant activity of coriander (*Coriandrum sativum* L.). *Acta Agrobotanica* 72 (1), 1–9.

Page AL, Miller RH, Keeney DR. 1982. *Method of Soil Analysis*. Part 2- Chemical and Microbiological Properties. Soil Science Society of America Inc., Madison, WI, USA.

Pratelli R, Pilot G. 2014. Regulation of amino acid metabolic enzymes and transporters in plants. *Journal of Experimental Botany* 65, 5535–5556.

Rather MA, Dar BA, Sofi SN. 2012. *Foeniculum vulgare*, a comprehensive review of its traditional use, phytochemistry, pharmacology, and safety. *Arabian Journal of Chemistry* 2, 1–10.

Saber M.S. 1997. Bio-fertilized Farming System. *Proceedings of the Training Course on Bio-organic Farming Systems for Sustainable Agriculture*, pp. 16–72.

Sadak M, Abdolhamid MT, Schmidhalter U. 2015. Effect of foliar application of amino acids on plant yield and some physiological parameters in bean plants irrigated with sea water. *Acta Biologica Colombiana* 20 (1), 141–152.

Schachtschable S. 1979. *Lehrbuch der Bodenkunde*. Printed in Germany (F.R.G.), Fedinaudenke, Stuttgart.

Shaji H, Chandran V, Mathew L. 2021. Chapter 13 -

Organic fertilizers as a route to controlled release of nutrients. In: Lewu FB, Volova T, Thomas S, KRR (eds) *Controlled Release Fertilizers for Sustainable Agriculture*. Academic Press, 231–245.

Shahhoseini R, Azizi M, Asili J, Moshtaghi N, Samiei L. 2020. Effects of zinc oxide nanoelicitors on yield, secondary metabolites, zinc and iron absorption of Feverfew (*Tanacetum parthenium* (L.) Schultz Bip.). *Acta Physiologiae Plantarum* 42, 1–18.

Shams M, Yildirim E, Ekinci M, Turan M, Dursun A, Parlakova F, Kul R. 2016. Exogenously applied glycine betaine regulates some chemical characteristics and antioxidative defense systems in lettuce under salt stress. *Horticulture, Environment and Biotechnology* 57 (3), 225–231.

Soliman YM, Abdul-Hafeez EY, Ibrahim OHM, Soliman TMA. 2023. Foliar application of glycine and/or zinc enhances vegetative, fruit and essential oil characters of *Cuminum cyminum* L. under different planting methods. *Assiut Journal of Agriculture Science* 54 (1), 66–84.

Souri MK, Hatamian M. 2019. Aminochelates in plant nutrition: A review. *Journal of Plant Nutrition* 42 (1), 67–78.

Taiwo L.B., Adediran J.A., Ashaye O.A., Odofin O., Oyadoyin A.J. 2002. Organic Okra (*Abelmoschus esculentus*): its growth, yield and organoleptic properties. *Nutrition & Food Science* 32(415), 180-183. (*Hort. Abst.* 72(12), 10918).

Taghizadeh SF, Azizi M, Rezaee R, Giesy JP, Karimi G. 2021. Polycyclic aromatic hydrocarbons, pesticides, and metals in olive: analysis and probabilistic risk assessment. *Environmental Science and Pollution Research* 28, 39723–39741.

Tao Y, Ferrer J, Ljung K, Pojer F, Hong F, Long J.A, Li L, Moreno J.E, Bowman M.E, Ivans L.J, Cheng Y, Lim J, Zhao Y, Ballaré C.L, Sandberg G, Noel J.P, Chory J. 2008. Rapid synthesis of auxin via a new tryptophan-dependent pathway is required for shade avoidance in plants. *Cell* 133 (1): 164–176.

Yassen AA, Mazher AAM, Zaghoul SM. 2010. Response of anise plants to nitrogen fertilizer and foliar spray of tryptophan under agricultural drainage water. *New York Science Journal* 3 (9), 120–127.